

EFFECT OF SOIL MOISTURE DEPLETION AND NITROGEN LEVELS ON WHEAT (*Triticum aestivum* L.).

El-Agrodi, M.W.M.¹; M.M. Saeid²; G. L. Ahmed¹ and T.S.H. Khalifa²

1- Soil Dept. Fac. of Agric., Mansoura Univ., Egypt.

2- Soil, Water and Environment Res. Inst., A.R.C., Egypt.



ABSTRACT

Lysimeter experiment was conducted during winter season of 2013-2014 and 2014-2015 in Sakha Agricultural Research Station Farm; Kafr El-Sheikh Governorate. This study aimed to clarify the effect of soil moisture depletion (45, 60 and 75% from the available water) and nitrogen levels (zero, 75, 100 and 125% from the recommended dose) as well as their interaction on wheat yield, water and nitrogen use efficiencies.

The obtained results can be summarized in;

- Soil moisture depletion and nitrogen fertilizer levels interaction high significantly affected grain yield in both seasons. The highest values of grain yield (2889.60 and 2829.40 kg fed⁻¹ in the 1st and 2nd season, respectively) were realized by applying the treatment of irrigation after depletion 45% from available water + 125% of N recommended dose.
- The highest values of wheat straw yield (3355.19 and 3297 kg fed⁻¹) were obtained under irrigation after depletion 45% from available water in the first and second season, respectively. Also the different N rates (control, 75%, 100% and 125% of N recommended dose) high significantly affected straw yield. The straw yield increased by 22.46% at N_{75%}, 31.61% at N_{100%} and 43.39% at N_{125%} in first season and by (18.71% at N_{75%}, 24.79% at N_{100%} and 26.26% at N_{125%}) in second season, as compared to control.
- The highest value of nitrogen agronomic efficiency (NAE) was obtained due to the irrigation after depletion 45% from available water + 75% of N recommended dose. The lowest value of NAE was achieved due to the irrigation after depletion 75% from available water + 125% of N recommended dose.
- Apparent nitrogen recovery by wheat was high significantly affected by the interaction between soil moisture depletion and nitrogen fertilizer levels in both seasons. The highest values of apparent nitrogen recovery by wheat were realized by applying the treatment of irrigation after depletion 45% from available water + 75% of N recommended dose for both seasons.
- The highest and lowest values of water use efficiency were obtained from treatments of irrigation after depletion 45% from available water and 75% from N recommended dose and irrigation after depletion 75% from available water without addition of nitrogen fertilizer in both growing seasons, respectively.

Keywords: Wheat, nitrogen agronomic efficiency, apparent nitrogen recovery, soil moisture depletion, field water use efficiency.

INTRODUCTION

The domestic wheat production in 2013 season was estimated by 8.7 million tons, whereas the Egyptian national consumption of wheat was about 17.7 million tons in 2013; there is a great gap between the consumption and production. Egypt imports above five million tons of wheat grains. Unless domestic wheat production increases annually, the deficit will increase due to the increase of population rate (2%) and present consumption per capita estimated by about 200 kg year⁻¹ (USDA, 2013).

Recently, water scarcity became an alarming problem. Water stress may result in similar yield or low yield reduction, with the benefit that the amount of saved irrigation water can be assigned to be used in irrigating new lands. However, water is an affecting factor in any agricultural expansion. Accordingly, it is advised to evaluate possible new approaches to control the crop water requirements through modern irrigation systems and management techniques. So, the use of improved irrigation systems becomes very important to save water, give favorable crop yield, optimum use of water and minimum labors requirements. Many researchers proved the importance of irrigation treatment to maximize wheat productivity. In this respect, Beshara (2012) stated that the highest value of the grain, straw, total N uptake and WUE were obtained with 40% depletion of available soil moisture in the growing season. While the lowest value of these traits were obtained with 70% depletion of available soil

moisture in both seasons. Hammad- Salwa and Ali (2014) indicated that delaying irrigation until 65% or 80% depletion decreased grain and straw yields. Increasing soil moisture depletion levels decreased the grain and straw yields compared with 50% depletion of the two seasons. They found also that total N uptake by wheat went as the same trend of grain and straw yields. Mahamed *et al.*, (2011) suggested that the greatest water use efficiency was observed when irrigation schedule applied when 50% of the available soil water is depleted as compared with Irrigation was applied when the soil moisture was depleted by 60% and 75% of available soil water (ASW).

Egypt utilizes fertilizers at an accelerating rate, due to various factors such as the increase in the cropped area and raising the rate of fertilizer application for various crops. Consequently, Egypt is considered to be a heavy user of chemical fertilizers, especially N fertilizers then P and K fertilizers. Soil fertility continues to decline because of combined effects of increasing pressure for land use for crop production, inadequate compensation of nutrients exported and lack of nutrients management. Gazia and Abd EL Aziz (2013) concluded that increasing N level up to 90 kg fed⁻¹ led to a significant increase in wheat grain yield. Mosaad *et al.*, (2013) showed that wheat grain and straw yields and N-uptake by grain and straw increased with use of mineral fertilizer rates up to 90 kg N fed⁻¹, but N utilization efficiency for wheat crop was decreased. Rahman *et al.*, (2011) reported that apparent nitrogen use efficiency (NUE) in terms of amount grain

production per unit N applied significantly decreased with the increasing N rate. Under the sub-optimum rate of 80 kg N ha⁻¹ application of all basal gave the maximum NUE, where minimum NUE was obtained under N rate of (120 kg ha⁻¹). Also they found that the highest values for total nitrogen uptake were achieved with N rate of (120 kg ha⁻¹). Noureldin *et al.*, (2013) reported that supplying wheat plants by 50 kg N fed⁻¹ produced the maximum grain yield/ N unit and the greatest apparent nitrogen use efficiency (AE). They also found that low apparent nitrogen use efficiency (AE) was achieved due to adding 100 kg N fed⁻¹. El-Gizawy (2005) indicated that grain yield fed⁻¹ was increased due to increasing the rates of N from 50 to 75 and 100 kg N fed⁻¹ in both season. Faizy *et al.*, (2010) stated that nitrogen use efficiency (NUE) and nitrogen recovery of wheat yield decreased with increasing N-level up to 140 kg fed⁻¹, where the highest value of NUE and nitrogen recovery obtained with 40 kg N fed⁻¹ as compared with 140 kg N fed⁻¹. Haile *et al.*, (2012) reported that the highest N uptake efficiencies were recorded at 30 kg N ha⁻¹ and the lowest N uptake efficiencies were recorded at the highest N rate (120 kg N ha⁻¹).

Therefore, the objective of this study is to evaluate the effect of soil moisture depletion levels and different nitrogen rates on wheat plant.

MATERIALS AND METHODS

Lysimeter experiment was conducted at Sakha Agricultural Research Station Farm; Kafr El-Sheikh Governorate, North-Nile Delta. The study aimed to clarify the effect of different soil moisture depletion, nitrogen fertilization and their interaction on wheat yield, water and nitrogen use efficiencies.

The studied factors were three soil moisture depletion (irrigation after depletion 45%, 60% and 75% from available water) and four nitrogen fertilization levels (0, 75, 100 and 125 % of nitrogen recommended dose of Agric. Ministry). The experiment was conducted in split plot design, with three replicates.

Table (1): The mean values of number of irrigation, total amount of water applied for each irrigation and the rainfall (R) for both seasons.

depletion	N-levels	Season 2013/2014			Season 2014/2015		
		No. irr.	Total Wa	R	No. irr.	Total Wa	R
45 %	N ₀	6	1105.11	173.43	6	1105.45	161.89
	N _{75%}	7	1298.15	173.43	7	1293.16	161.89
	N _{100%}	8	1570.96	173.43	8	1576.05	161.89
	N _{125%}	9	1819.94	173.43	9	1817.21	161.89
60 %	N ₀	4	1053.84	173.43	4	1062.44	161.89
	N _{75%}	5	1234.07	173.43	5	1248.20	161.89
	N _{100%}	6	1445.32	173.43	6	1482.80	161.89
	N _{125%}	7	1618.41	173.43	7	1715.61	161.89
75 %	N ₀	3	988.49	173.43	3	998.66	161.89
	N _{75%}	4	1174.79	173.43	4	1168.40	161.89
	N _{100%}	5	1398.73	173.43	5	1390.58	161.89
	N _{125%}	6	1573.90	173.43	6	1666.93	161.89

Soil samples (0-15 , 15-30 and 30-45 cm depth) from each lysimeter were taken before planting and

Lysimeter (0.78 m²) was divided into 3 groups; each group includes 12 lysimeters to be studied.

Lysimeter was a circular shape; the diameter of one metre and a height of 60 cm with filter (sand and gravel) of 10 cm, each lysimeter was filled by 458.25 kg of the clay soil. The area of Lysimeter was determined using the formula: Area = π × r². Nitrogen recommendation for wheat 75 kg N fed⁻¹. Urea fertilizer was used as a source of nitrogen

Wheat (*Triticum aestivum L.*) variety (*Sakha 93*) was sown in first season on December, 5th, 2013 and the second season was sown on December, 3rd, 2014. Harvesting was done on May, 20th, 2014 and on May, 18th, 2015 after full maturity for both seasons. The wheat was sown at the rate of 60 kg of seeds fed⁻¹. Each of N dose was divided into two equal applications; one was added after 25 days from sowing and the second at the following irrigation. Other agricultural practices such as weed and insect control were performed according to the Ministry of Agriculture recommendation in North Nile Delta area.

The irrigation water applied was calculated to raise the soil water content of the upper 15 cm to field capacity in the first month, of the upper 30 cm in the second month and of the upper 45 cm in the rest period. Irrigation was stopped three weeks before harvesting.

Actual water consumptive use consists of the two components;

1-water applied (Wa) where water applied for irrigation was determined according to Phocaides (2001) as follow:-

$$(Wa) = f (Fc - WP) * BD * Ds$$

Where:

Fc = field capacity (%), WP = wilting point (%).
 f = permissible depletion, BD = bulk density (g cm⁻³)
 Ds = soil layer

2- rainfall (R), where the total amount of the rainfall (R) was 41.29 mm (173.43 m³ fed⁻¹) and 38.55 mm (161.89 m³ fed⁻¹) in the 1st and 2nd seasons, respectively. The number of irrigation, total amount of water applied for each irrigation and the rainfall (R) are shown in Table (1).

Table (1): The mean values of number of irrigation, total amount of water applied for each irrigation and the rainfall (R) for both seasons.

depletion	N-levels	Season 2013/2014			Season 2014/2015		
		No. irr.	Total Wa	R	No. irr.	Total Wa	R
45 %	N ₀	6	1105.11	173.43	6	1105.45	161.89
	N _{75%}	7	1298.15	173.43	7	1293.16	161.89
	N _{100%}	8	1570.96	173.43	8	1576.05	161.89
	N _{125%}	9	1819.94	173.43	9	1817.21	161.89
60 %	N ₀	4	1053.84	173.43	4	1062.44	161.89
	N _{75%}	5	1234.07	173.43	5	1248.20	161.89
	N _{100%}	6	1445.32	173.43	6	1482.80	161.89
	N _{125%}	7	1618.41	173.43	7	1715.61	161.89
75 %	N ₀	3	988.49	173.43	3	998.66	161.89
	N _{75%}	4	1174.79	173.43	4	1168.40	161.89
	N _{100%}	5	1398.73	173.43	5	1390.58	161.89
	N _{125%}	6	1573.90	173.43	6	1666.93	161.89

after harvesting in the two seasons, air-dried, gently crushed and sieved to pass a 2 mm sieve to determine

some chemical and physical soil properties as shown in Table (2).

Method of analysis:

- Soil reaction (pH) according to Cottenie *et al.*, (1982), Electrical conductivity (ECe), soluble cations and anions according to Page (1982). Organic matter (OM) content was determined using (Walkally Black - method) according to Hesse, (1971). Available N and K were determined according to Jackson (1967). Available P was determined by sodium bicarbonate

method according to Olsen *et al.*, (1954). The cation exchange capacity (CEC) was determined as described by Jackson (1967). Mechanical soil analysis and CaCO₃% were determined according to the international pipette method (Piper, 1950). Field capacity (F.C.) and permanent wilting point (PWP) were determined by using pressure membrane method at 0.33 and 15 atm (Klute, 1986). Available water value is the difference between the field capacity and the permanent wilting point.

Table (2): The average values of some chemical, physical and soil moisture characteristics of the experimental soil before wheat planting.

Seasons		1 st	2 nd
pH 1:2.5		8.33	8.28
EC _e dS/m		2.90	2.73
Soluble cations (meq /L)	Na ⁺	18.99	17.83
	Ca ⁺⁺	4.06	3.86
	Mg ⁺⁺	6.03	5.33
	K ⁺	0.25	0.27
Soluble anions (meq /L)	CO ₃ ⁻⁻	0	0
	HCO ₃ ⁻	3.33	3.17
	Cl ⁻	14.04	13.46
	SO ₄ ⁻⁻	11.95	10.67
Available (mg/kg)	N	17.17	19.41
	P	9.18	12.47
	K	303.43	290.42
Soil mechanical analysis (%)	Sand	18.61	20.11
	Silt	31.68	31.99
	Clay	49.71	47.9
Texture grade		clay	clay
OM %		1.15	1.10
Total CaCO ₃ (%)		2.32	2.28
CEC meg/100 g soil		38.58	39.50
Field capacity (%)		39.26	39.38
Wilting point (%)		21.34	21.40
Available water (%)		17.92	17.98
Bulk density (g cm ⁻³)		1.17	1.18

Dry materials of plant organs (oven dry basis) were wet digested with an H₂SO₄ – H₂O₂ mixture as described by Wolf (1982) and analyzed for the following:

- Nitrogen content was determined by micro-Kjeldahl method (Jackson, 1967).
- Phosphorus was determined by using hydroquinine method (Snell and Snell, 1967).
- Potassium content was determined by using a flame photometer (Jackson 1967).

The following parameters were calculated:

- **Apparent N Recovery (ANR):** by the equation described by Echeverria and Videla (1998), *i.e.*, ANR (%) = [N uptake (fertilized plot) – N uptake (zero plot) / N fertilizer rate] X 100.
- **Nitrogen Agronomic Efficiency (NAE) according to Craswell and Godwin (1984):** [grain yield (fertilized plot) - grain yield (zero plot)] / N rate.
- **Water use efficiency (WUE):** was calculated by dividing the total weight of grain yield (kg fed⁻¹) by the amount of seasonal water consumptive use (m³ fed⁻¹) Talha *et al.*, (1980).

$$WUE (Kg m^3) = \frac{Grain\ yield\ (kg\ fed^{-1})}{WCU\ (m^3\ fed^{-1})}$$

Where: WCU = actual water consumptive use

All measurements data during the two seasons in this study were statistically analyzed according to the methods described by Snedecor and Cochran (1980). The differences among the means of different treatments were tested using the Least Significant Differences (LSD) at probability 5%. Statistical analysis was done using the CoStat package program, version 6.311 (Cohort software, USA).

RESULTS AND DISCUSSION

1- Grain yield:

Data presented in Table (3) show the effect of soil moisture depletion, nitrogen fertilization levels and their interaction on grain yield.

Data point out that soil moisture depletion treatment high significantly decreased the grain yield in both seasons. Means of wheat grain yield were 2157.31, 2006.46, 1807.58 kg fed⁻¹ and 2219.35, 1951.60, 1825.25 kg fed⁻¹ under irrigation after depletion 45%,

60% and 75% from available water in the first and second season, respectively. Hence irrigation after depletion 45% from available water increase grain yield by (19.35% and 21.59%) than that irrigation after depletion 75% from available water in the first and second season, respectively. This is to be expected since drought stress might reduce translocation of assimilates from leaves, this response in addition to reduce photosynthesis in the leaves itself contribute to lower grain yield. Conversely, water deficit at any growth stage reduce grain yield. These results are in the same line with Beshara (2012) and Hammad - Salwa and Ali (2014).

Adding nitrogen fertilizer up to 125% from the recommended dose (93.75 kg N fed⁻¹) high significantly increased grain yield (kg fed⁻¹). Nitrogen application at rate of 125% increases grain yield by 1647.42, 585.95 and 282.43 kg fed⁻¹ in the first season and by 1629.6, 412.07 and 190.87 kg fed⁻¹ in the second season than that of control, 75% and 100% of N recommended dose, respectively. It can be concluded that the increasing

amount of nitrogen fertilizer, consequently get higher grain yield. These results are in conformity with El-Gizawy (2005); Mosaad *et al.*, (2013) and Gazia and Abd EL Aziz (2013).

High significantly effect was found by the interaction between soil moisture depletion and nitrogen fertilizer levels on grain yield in both seasons. The highest values of grain yield are realized by applying the treatment of irrigation after depletion 45% from available water + 125% of N recommended dose for both seasons. Grain yield reduced in each season by increasing soil moisture depletion at all nitrogen rates in both seasons. Data of Table 3 state also that the obtained grain yield from applying the recommended N dose (75 kg N fed⁻¹) and irrigation after depletion 45% from available water (2542.40 kg fed⁻¹ in the 1st season) can be realized by increasing the dose of N to 125% from the recommended (93.75 kg N fed⁻¹) and delay the irrigation to be done after depletion 60% from available water (2633.40 kg grains fed⁻¹ in the 1st season)

Table (3): Effect of soil moisture depletion, N fertilization levels and their interactions on grain yield (kg fed⁻¹)

depletion	Grain yield (kg fed ⁻¹)				F-test
	45%	60%	75%	Mean	
Seasons					
2013 - 2014					
N-levels					**
N ₀	1046.85	989.45	879.64	971.98	LSD
N _{75%}	2210.60	2056.60	1833.16	2033.45	0.05= 31.48
N _{100%}	2542.40	2346.40	2122.12	2336.97	
N _{125%}	2829.40	2633.40	2395.40	2619.40	0.01= 43.13
Mean	2157.31	2006.46	1807.58		
F-test	**	LSD 0.05= 48.15	0.01= 79.84		
Interaction					
2014 - 2015					
N-levels					**
N ₀	988.40	935.20	858.20	927.27	LSD
N _{75%}	2322.60	2118.20	1993.60	2144.80	0.05=51.75
N _{100%}	2676.80	2294.60	2126.60	2366.00	
N _{125%}	2889.60	2458.40	2322.60	2556.87	
Mean	2219.35	1951.60	1825.25		0.01=70.90
F-test	**	LSD 0.05=74.06	0.01=122.81		
Interaction					

2-Straw yield:

Data presented in Table (4) show the effect of soil moisture depletion, nitrogen fertilization levels and their interaction on wheat straw yield.

Data indicate that soil moisture depletion treatments significantly decreased wheat straw yield in both seasons. The highest mean values of wheat straw yield are 3355.19 and 3297 kg fed⁻¹ under irrigation after depletion 45% from available water in the first and second season, respectively. No significant differences in the straw yield were found as affected by irrigation after depletion 45% and 60% from available water in first season. Also the lowest mean values are 3080.98 and 3127.25 kg fed⁻¹ under irrigation after depletion 75% from available water in the first and second season, respectively, without significant difference as compared with irrigation after depletion 60% from available water

in the second season. The increase in growth and productivity of wheat due to increasing irrigation water may be due to provide moisture for wheat plants continuously which allows better growth, thereby enhancement yield components resulting in increments in plant height and straw yield. Similar results are given by previous researchers Beshara, (2012) and Hammad-Salwa and Ali (2014)

Different N rates 75%, 100% and 125% of N recommended dose high significantly increased straw yield by 22.46% at N_{75%}, 31.61% at N_{100%} and 43.39% at N_{125%} in first season and by 18.71% at N_{75%}, 24.79% at N_{100%} and 26.26% at N_{125%} in second season, as compared to control. Insignificant differences between 100% and 125% of N recommended dose in second season were found. It can be concluded that the increasing amount of nitrogen fertilizer, the straw yield

gradually improve because encourage vegetative growth and thus on plant height and branches, so on the straw yield. These results are in harmony with those obtained by Mosaad *et al.*, (2013).

No interaction effects were found between soil moisture depletion and nitrogen fertilization levels on wheat straw yield in both seasons.

Table (4): Effect of soil moisture depletion, fertilization levels and their interactions on straw yield (kg fed⁻¹):

depletion	Straw yield (kg fed ⁻¹)				F-test
	45%	60%	75%	Mean	
Seasons		2013 - 2014			
N-levels					**
N ₀	2754.15	2677.85	2410.36	2614.12	LSD
N _{75%}	3270.40	3249.40	3084.06	3201.29	0.05=72.75
N _{100%}	3540.60	3505.60	3274.88	3440.36	
N _{125%}	3855.60	3834.60	3554.60	3748.27	0.01=99.67
Mean	3355.19	3316.86	3080.98		
F-test	**	LSD 0.05=79.38	0.01=131.64		
Interaction					ns
Seasons		2014-2015			
N-levels					**
N ₀	2840.60	2711.80	2620.80	2724.40	LSD
N _{75%}	3320.80	3201.80	3179.40	3234.00	0.05=86.68
N _{100%}	3469.20	3382.40	3347.40	3399.67	
N _{125%}	3557.40	3400.60	3361.40	3439.80	0.01=118.76
Mean	3297.00	3174.15	3127.25		
F-test	*	LSD 0.05=83.76	0.01=138.89		
Interaction					ns

3-Total Nitrogen uptake by wheat:

Data presented in Table (5) show the effect of soil moisture depletion, nitrogen fertilization levels and their interactions on total nitrogen uptake by wheat (grain + straw uptake).

Total nitrogen uptakes are high significantly affected by soil moisture depletion treatments in the two growing seasons. Mean values are 54.32, 49.34, 43.38 kg N fed⁻¹ and 62.22, 53.86, 46.97 kg N fed⁻¹ under irrigation after depletion 45%, 60% and 75% from available water in first and second seasons, respectively.

Irrigation after depletion 45% from available water increased the total N uptake by 4.98, 10.94 kg N fed⁻¹ and by 8.36, 15.25 kg N fed⁻¹ in the 1st and 2nd season as compared with irrigation after depletion 60% and 75% from available water, respectively. These results may be attributed to the increase in soil moisture content in the plant root zone which led to an increase in available nutrient and increase N uptake by wheat plant. These results were recorded by Beshara, (2012) and Hammad - Salwa and Ali (2014).

Table (5) Effect of soil moisture depletion and nitrogen fertilization levels on total N uptake (kg N fed⁻¹):

depletion	Total N uptake (kg N fed ⁻¹)				F-test
	45%	60%	75%	Mean	
seasons		2013 - 2014			
N-levels					**
N ₀	21.90	19.30	16.38	19.19	LSD
N _{75%}	55.17	50.18	44.87	50.07	0.05=1.57
N _{100%}	65.44	59.35	51.99	58.93	
N _{125%}	74.77	68.53	60.29	67.86	0.01=2.16
Mean	54.32	49.34	43.38		
F-test	**	LSD 0.05=1.07	0.01=1.77		
Interaction					**
seasons		2014 - 2015			
N-levels					**
N ₀	24.16	19.64	15.96	19.92	LSD
N _{75%}	65.30	56.20	49.71	57.07	0.05=1.75
N _{100%}	75.18	65.65	57.02	65.95	
N _{125%}	84.24	73.96	65.18	74.46	0.01=2.40
Mean	62.22	53.86	46.97		
F-test	**	LSD 0.05=2.17	0.01=3.59		
Interaction					**

Data in Table 5 also reveal that nitrogen fertilization levels in both seasons high significantly affected the total N uptake by wheat plant. The highest values of total nitrogen uptake are 67.86 and 74.46 kg N fed⁻¹ under 125% of N recommended dose while the lowest values are 19.19 and 19.92 kg N fed⁻¹ under control (zero N addition. Mean values of 75% and 100% of N recommended dose are 50.07, 58.93 kg N fed⁻¹ and 57.07, 65.95 kg N fed⁻¹ in 1st and 2nd seasons, respectively. This may be due to the available nitrogen in the soil are less under the low levels and are enough in the high level, also increasing nitrogen fertilizer levels led to increase nitrogen uptake by plants. These results are in line with Rahman *et al.*, (2011).

There are high significant differences in total nitrogen uptake in the both seasons due to the interaction between soil moisture depletion and nitrogen fertilization levels. It could be noticed that irrigation after depletion 45 % from available water with fertilization at 125 % of recommended dose gave the highest values of nitrogen total uptake by wheat. On the contrary, the lowest total nitrogen uptakes by wheat are

obtained from irrigation at 75 % available water without nitrogen fertilization in the two growing seasons.

4- Nitrogen agronomic efficiency (NAE):

Data presented in Table (6) show the effect of soil moisture depletion, nitrogen fertilization levels and their interaction on nitrogen agronomic efficiency by wheat grains (kg grain kg N⁻¹).

Data reveal that soil moisture depletion significantly affected NAE, where irrigation after depletion 75 % from available water decreased NAE by 16.70 % and 9.01 % than that of irrigation after depletion 45 % and 60% from available water, in first season, respectively. While in second season these decreases were 20.74 % and 4.87% than that of irrigation after depletion 45 % and 60% from available water, respectively. Insignificant differences were found between mean values of NAE due to irrigation after depletion 60 % and 75% from available water in second season. The data reveal that the unit of nitrogen (kg) produces more grains yield when plant was irrigated after depletion of 45% from available water than 60% and 75%.

Table (6) Effect of soil moisture depletion and nitrogen fertilization levels on nitrogen agronomic efficiency (NAE) Kg grains Kg N⁻¹ for wheat:

depletion	NAE (Kg grains Kg N ⁻¹)				F-test
	45%	60%	75%	Mean	
seasons					
2013 - 2014					
N-levels					**
N ₀	-	-	-	-	LSD
N _{75%}	20.69	18.97	16.95	18.87	0.05= 0.41
N _{100%}	19.94	18.09	16.57	18.20	0.01= 0.57
N _{125%}	19.01	17.53	16.17	17.57	
Mean	19.88	18.20	16.56		
F-test	**	LSD 0.05=0.66	0.01=1.09		
Interaction					**
seasons					
2014-2015					
N-levels					**
N ₀	-	-	-	-	LSD
N _{75%}	23.72	21.03	20.18	21.93	0.05= 0.67
N _{100%}	22.51	18.13	16.91	19.59	0.01= 0.91
N _{125%}	20.28	16.25	15.62	17.59	
Mean	22.17	18.47	17.57		
F-test	**	LSD 0.05=1.27	0.01=2.11		
Interaction					**

Nitrogen fertilization levels greatly affect NAE, where significant decreases were found with increasing N fertilization level. The highest mean values 18.87 and 21.93 kg grain kg N⁻¹ were achieved under application of 75% of N recommended dose and the lowest mean values 17.57 and 17.59 kg grain kg N⁻¹ were recorded under application of 125% of N recommended dose. While mean values were 18.20 and 19.59 kg grain kg N⁻¹ under application of 100% of N recommended dose in the 1st and 2nd seasons, respectively. These results are in agreement with Rahman *et al.*, (2011) and Nouredin *et al.*, (2013).

Concerning the interaction effect data reveal that high significant interactions between soil moisture depletion and N fertilization levels on NAE were obtained. The highest values of NAE 20.69 and 23.72

kg grain kg N⁻¹ were recorded under irrigation after depletion 45% from available water + 75% of N recommended dose. The lowest values (16.17 and 15.62 kg grain kg N⁻¹) were achieved under irrigation after depletion 75% from available water + 125% of N recommended dose. It can be said that NAE increases by decreasing soil moisture depletion and N-levels in both seasons.

5- Apparent N recovery (ANR) by wheat (%):

Data presented in Table (7) show the effect of soil moisture depletion, nitrogen fertilization levels and their interaction on apparent nitrogen recovery (ANR) by wheat (%).

Data point out that soil moisture depletion treatment high significantly affected ANR in both seasons. The mean values of apparent nitrogen recovery

by wheat are 57.87, 53.61, 48.13 % and 68.41, 61.43 and 55.74 % under irrigation after depletion 45%, 60% and 75% from available water in the first and second season, respectively. Hence irrigation after depletion 75% from available water decreased ANR by (17.62 % and 10.55 %) and (19.46 % and 10.06 %) than that irrigation after depletion 45% and 60% from available water in the first and second season, respectively.

Raising nitrogen fertilizer levels negatively and high significantly affected ANR (%). 125% of N recommended dose decreased ANR by (5.38 and 1.98 %) in the first season and (11.92 and 5.21 %) in the second season than that of 75% and 100% of N

recommended dose, respectively. Similar results were given by previous researchers Faizy *et al.*, (2010) and Haile *et al.*, (2012).

Apparent nitrogen recovery values by wheat were high significant affected by the interaction between soil moisture depletion and nitrogen fertilizer levels in both seasons. The highest values of apparent nitrogen recovery by wheat were realized by applying the treatment of irrigation after depletion 45% from available water + 75% of N recommended dose for both seasons. Apparent nitrogen recovery by wheat reduced in each season by increasing soil moisture depletion at all nitrogen rates in both seasons.

Table (7) Effect of soil moisture depletion and N fertilization levels on apparent N recovery (ANE) % by wheat (grains + straw):

depletion	Apparent N recovery (%)				F-test
	45%	60%	75%	Mean	
Seasons		2013 - 2014			**
N-levels					LSD
N ₀	-	-	-	-	0.05=2.08
N _{75%}	59.14	54.91	50.41	54.82	
N _{100%}	58.06	53.40	47.30	52.92	0.01=2.85
N _{125%}	56.40	52.51	46.69	51.87	
Mean	57.87	53.61	48.13		
F-test	**	LSD 0.05= 2.29	0.01= 3.79		
Interaction					**
Seasons		2014-2015			**
N-levels					LSD
N ₀	-	-	-	-	0.05=2.50
N _{75%}	73.13	65.01	60.00	66.04	
N _{100%}	68.02	61.35	54.74	61.37	0.01=3.43
N _{125%}	64.08	57.94	52.49	58.17	
Mean	68.41	61.43	55.74		
F-test	**	LSD 0.05= 3.25	0.01=5.40		
Interaction					**

6- Actual water consumptive use:

Data presented in Table (8) show the effect of soil moisture depletion, nitrogen fertilization levels and

their interaction on actual water consumptive use of wheat yield.

Table (8) Effect of soil moisture depletion and N fertilization levels on actual water consumptive use (m³ fed⁻¹):

depletion	Actual water consumptive use (m ³ fed ⁻¹)			
	45%	60%	75%	Mean
Seasons		2013 - 2014		
N-levels				
N ₀	1278.54	1227.27	1161.92	1222.58
N _{75%}	1471.58	1407.5	1348.22	1409.10
N _{100%}	1744.39	1618.75	1572.16	1645.10
N _{125%}	1993.37	1799.33	1747.33	1846.68
Mean	1621.97	1513.21	1457.41	
Seasons		2014-2015		
N-levels				
N ₀	1267.34	1224.33	1160.55	1217.41
N _{75%}	1455.05	1410.09	1330.29	1398.48
N _{100%}	1737.94	1644.69	1552.47	1645.03
N _{125%}	1979.10	1877.50	1828.82	1895.14
Mean	1609.86	1539.15	1468.03	

Data state that the decreasing % in actual water consumptive use values under irrigation after depletion 60% and 75% of available water compared to irrigation

after depletion 45% of available water by (6.71 and 10.15%) and (4.39 and 8.81%) in 1st and 2nd seasons respectively. These results are attributed to more

available soil moisture, under irrigation after depletion 45% of available water, which enhanced both transpiration from plants leaves and evaporation from the soil surface.

The addition of 125% from recommended dose cause increasing in actual water consumptive use of wheat crop. Average values of actual water consumptive use are (1846.68 and 1895.14 m³ fed⁻¹), (1645.10 and 1645.03 m³ fed⁻¹) and (1409.10 and 1398.48 m³ fed⁻¹) for 125%, 100% and 75% from recommended N dose in both seasons, respectively, these results could be caused due to increase the wheat yield which led to increases in water consumptive use.

Concerning to the interaction effect between soil moisture depletion and nitrogen fertilizer levels effect on actual water consumptive use in both seasons, data reveal that the highest values were obtained with irrigation at depletion of 45% of available water + 125%

of N recommended dose comparing with other soil moistures and nitrogen rates.

7- Water use efficiency:

Data presented in Table (9) show the effect of soil moisture depletion, nitrogen fertilization levels and their interaction on water use efficiency.

Water use efficiency for the grain yield was affected by soil moisture depletion levels and was highly significant. The mean values of WUE due to the irrigation after depletion 45% of available water are significantly different from that irrigation after depletion 60% and 75% of available water. The highest WUE (1.31 and 1.34 kg grain m⁻³ water consumed) and the lowest (1.21 and 1.23 kg grain m⁻³ water consumed) for both seasons were observed at 45 and 75 % SMD levels, respectively. These indicated that 75 % of soil moisture depletion reduce WUE for grain yield. The results agree with the result of Mahamed *et al.*, (2011) and Beshara (2012).

Table (9) Effect of soil moisture depletion and N fertilization levels on water use efficiency (kg grain m⁻³ water consumed)

depletion	WUE (kg grain m ⁻³ water consumed)				F-test
	45%	60%	75%	Mean	
Seasons		2013 - 2014			**
N-levels					LSD
N ₀	0.83	0.81	0.76	0.80	0.05= 0.02
N _{75%}	1.52	1.46	1.38	1.45	
N _{100%}	1.46	1.43	1.37	1.42	0.01= 0.03
N _{125%}	1.43	1.40	1.31	1.38	
Mean	1.31	1.28	1.21		
F-test	** LSD		0.05= 0.03	0.01= 0.05	
Interaction		ns			
Seasons		2014-2015			**
N-levels					LSD
N ₀	0.77	0.76	0.74	0.76	0.05= 0.03
N _{75%}	1.58	1.51	1.48	1.52	
N _{100%}	1.54	1.42	1.35	1.44	0.01= 0.04
N _{125%}	1.45	1.37	1.33	1.38	
Mean	1.34	1.27	1.23		
F-test	** LSD		0.05= 0.02	0.01= 0.04	
Interaction		*			

The effect of nitrogen at various levels on WUE were high statistically different in all cases. Maximum WUE of (1.45 and 1.52 kg grain m⁻³ water consumed) were obtained when N apply at level of 75% from N recommended dose. The lowest WUE (0.80 and 0.76 kg grain m⁻³ water consumed) were achieved without nitrogen application for both seasons, respectively.

Likewise, the interactions between soil moisture depletion and nitrogen fertilizer levels on WUE were insignificant in first season and significant in second seasons. The highest value of WUE was obtained from treatments of irrigation after depletion 45% of available water and 75% from N recommended dose while the lowest value of WUE was revealed from the irrigation after depletion 75% of available water without addition of nitrogen fertilizer in both growing seasons.

These results could be attributed to the highly significant differences between the wheat grain yield values as well as differences water consumed use (Table 3 and 8).

CONCLUSION

It can be concluded that under natural conditions (water availability) wheat can be irrigated after depletion 45 % from available water with fertilization at 125 % of recommended dose which gave the highest mean values of grain production of wheat, On the other hand, under drought conditions (reducing water resources), other soil moisture treatments with different nitrogen levels can be used with expected decreasing in wheat productivity.

REFERENCES

- Beshara, A. T. (2012). Effect of Soil Moisture Depletion and Nitrogen Fertilizer Application Date on Wheat Yield, Water and Fertilizer Use Efficiencies in North Africa. Ph. D. Thesis, Natural Resources, I.A.R.S., Cairo Univ., Egypt.
- Cottenie, A.; Verloo, M.; Velghe, G and Kiekens, L. (1982). Biological and analytical aspects of soil pollution. Lab. Of Analytical Agro. State Univ. Gent-Belgium.
- Craswell, E. T. and D. C. Godwin, (1984). The efficiency of nitrogen fertilizers applied to cereals in different climates. Adv. In Plant Nutrition, New York, 1: 1-55.
- Echeverria, H. E. and C. C. Videla, (1998). Eficiencia fisiologica y de utilizacion de nitrogeno en trigo en la region pampeana Argentina. Ciencia del suelo 16: 83 – 87.
- El-Gizawy, N. Kh. B. (2005). Yield and nitrogen use efficiency as influenced by rates and sources of nitrogen fertilizers of some wheat varieties. Conference of Agronomy, Agron. Dept., Fac. Agric., Assiut Univ., 9:15-16.
- Faizy, S. E. D. A.; M. M. Rizk; M. M. Ragab and M. M. A. Amer, (2010). Response of wheat yield and apparent nitrogen recovery of fertilizer to mineral nitrogen and biofertilizer application in salt affected soils. J. Agric. Res. Kafr El-Sheikh Univ., 36 (1).
- Gazia, E. A. E. and M. A. Abd EL-Aziz, (2013). Maximizing wheat yield under N, K and B fertilization . J. Soil Sci. and Agric. Eng., Mansoura Univ., 4 (10): 1073 – 1084.
- Haile, D.; D. Nigussie and A. Ayana, (2012). Nitrogen use efficiency of bread wheat: Effects of nitrogen rate and time of application. J. Soil Sci. and Plant Nutr. 12 (3): 389-409.
- Hammad - Salwa. A. R. and O. A. M. Ali (2014). Physiological and biochemical studies on drought tolerance of wheat plants by application of amino acids and yeast extract. Ann. Agric. Sci., 59, (1): 133–145.
- Hesse, P. R. (1971). A Text book of Soil Chemical Analysis. John Murray Ltd, London, pp. 520.
- Jackson, M. L. (1967). "Soil Chemical Analysis". Prentice-Hall of India, New Delhi.
- Klute , A. (1986). Methods of soil analysis (part 1). Amer. soc. Of Agron., Inc. Madison, Wisconsin, USA. 3rd edition.
- Mahamed, M. B.; E. Sarobol; T. Hordofa; S. Kaewrueng and J. Verawudh, (2011). Effects of Soil Moisture Depletion at Different Growth Stages on Yield and Water Use Efficiency of Bread Wheat Grown in Semi-Arid Conditions in Ethiopia. Kasetsart J. (Nat. Sci.) 45: 201 – 208.
- Mosaad, I. S. M.; E. E. E. Khafagy and R. A. El-Dissoky, (2013). Effect of mineral, bio and organic nitrogen fertilization on wheat yield and nitrogen utilization efficiency and uptake at northern delta of Egypt. J. Soil Sci. and Agric. Eng., Mansoura Univ., 4 (10): 1101 – 1116.
- Noureldin Nemat, A.; H. S. Saady; F. Ashmawy and H. M. Saed, (2013). Grain yield response index of bread wheat cultivars as influenced by nitrogen levels. Annals of Agricultural Science 58(2), 147–152.
- Olsen, S. R.; C.V. Cale; F. S. Watenable and L. A. Dean, (1954). Estimation of available phosphorus in soil by extraction with sodium bicarbonate U.S. Dept., Agric. Circ., 939.
- Page A. L. (ed.) (1982). Methods of Soil Analysis, part 2: Chemical and Microbiological properties, (2nd ed.) American Society at Agronomy, Inc. Soil. Sci Soc. Of Am. Inc., Madison, Wisconsin, U S A.
- Phocaides , A. (2001). Handbook on pressurized irrigation techniques .Food and Agriculture Organization of the United Nations, Rome.
- Piper, C.S. (1950). Soil and Plant Analysis. Inter science Publication. New York.
- Rahman, M. A.; M. A. Z. Sarker; M. F. Amin; A. H. S. Jahan and M. M. Akhter, (2011). Yield response and nitrogen use efficiency of wheat under different doses and split application of nitrogen fertilizer. Bangladesh J. Agril. Res. 36(2): 231-240.
- Snedecor, G. W. and W. G. Cochran, (1980). Statistical Methods, seventh ed. The Iowa State Univ. Press, Ames. Iowa, USA, pp. 1–507.
- Snell, F. D. and C. T. Snell, (1967). Colorimetric Methods of Analysis. D. Van. Nostrand company Inc.: 551 – 552.
- Talha, M.; M. A. Aziz, and E. M. El-Toni, (1980). The combined effect of irrigation intervals and cycocel treatments on Pelargonium graveolens L. II- Evapotranspiration and water economy. Egypt. J. Soil Sci., 20(2): 121.
- USDA, (2013). USDA Gain: Egypt grain and feed annual 2013: Forex availability impacts grain imports. USDA Foreign Agricultural Service
- Wolf, (1982). A comprehensive system of leaf analysis and it is use for diagnosing crop nutrient status. Commun Soil Sci. and plant Analysis. 13: 1035-1059.

تأثير استنزاف الرطوبة الارضية و مستويات التسميد النيتروجيني على القمح.
محمد وجدى محمد العجرودى¹ ، محمود محمد سعيد²، جمعة لبيب احمد يوسف¹ و تامر حسن حلمى خليفة²
1- قسم الأراضى- كلية الزراعة- جامعة المنصورة
2- معهد بحوث الأراضى والمياه والبيئة - مركز البحوث الزراعية

أجريت تجربة ليزمبترات على محصول القمح شتاء عام 2014/2013 و 2015/2014 بالمزرعة البحثية بمحطة بحوث سخا- محافظة كفر الشيخ لتقييم أثر الرطوبة الارضية (الرى للسعة الحقلية بعد استنفاد (45، 60، 75 % من الماء الميسر) ومستويات التسميد النيتروجيني (صفر ، 75 ، 100 ، 125% من الموصى به (75 كجم نيتروجين / فدان) على محصول القمح والممتص من النيتروجين وكفاءة الاستخدام للماء و النيتروجين.

وفيما يلي أهم النتائج المتحصل عليها:

- أثرت الرطوبة الارضية ومستويات التسميد النيتروجيني والتفاعل بينهما تأثيرا عالى المعنوية على محصول الحبوب فى كلا الموسمين و سجلت اعلى قيمة له (2889.60، 2829.40 كجم / فدان) مع تطبيق الري بعد استنزاف 45% من الماء الميسر + 125 % من الموصى به للسماذ النيتروجيني فى كلا الموسمين.
- سجلت اعلى قيمة لمحصول القش (3355.19، 3297 كجم / فدان) مع تطبيق الري بعد استنزاف 45% من الماء الميسر فى الموسم الاول و الثانى على التوالي. وايضا مع معدلات النيتروجين المختلفة (75% ، 100% ، 125% من الموصى به) كانت لها تأثيرا عالى المعنوية على زيادة محصول القش بنسبة 22.46% ، 31.61% ، 43.39% بالموسم الاول و بنسبة 18.71% ، 24.79% ، 26.26% بالموسم الثانى مع معدلات النيتروجين المختلفة (75% ، 100% ، 125% من الموصى به) على التوالي مقارنة بالكنترول.
- سجلت اعلى قيمة لكفاءة النيتروجين المحصولية (NAE) (20.69 ، 23.72 كجم حبوب / كجم نيتروجين مضاف) مع تطبيق الري بعد استنزاف 45% من الماء الميسر + 75 % من الموصى به للسماذ النيتروجيني. بينما اقل قيمة (16.17 ، 15.62 كجم حبوب / كجم نيتروجين مضاف) مع تطبيق الري بعد استنزاف 75% من الماء الميسر + 125 % من الموصى به للسماذ النيتروجيني فى كلا الموسمين.
- التفاعل بين استنزاف الرطوبة الارضية ومستويات التسميد النيتروجيني كان لها تأثيرا عالى المعنوية على الكمية الممتصة من النيتروجين (ANR) فى كلا الموسمين. و سجلت اعلى قيمة للكمية الممتصة من النيتروجين مع تطبيق الري بعد استنزاف 45% من الماء الميسر + 75 % من الموصى به للسماذ النيتروجيني فى كلا الموسمين.
- سجلت اعلى قيمة لكفاءة استخدام القمح للمياه مع تطبيق الري بعد استنزاف 45% من الماء الميسر + 75 % من الموصى به للسماذ النيتروجيني و اقل قيمة مع تطبيق الري بعد استنزاف 75% من الماء الميسر بدون اضافة للسماذ النيتروجيني لكلا الموسمين على التوالي.