

## **Allivating Salinity and Sodicy by Adding Some Soil Amendments**

**Gihan H. Kamel<sup>1,2</sup>; Esmat Noufal<sup>2</sup>; I. Farid<sup>2</sup>; S. Abdel-Aziz<sup>1</sup> and M. H. H. Abbas<sup>1,\*</sup>**

<sup>1</sup> Soils, Water and Environment Research Institute (SWERI), Agricultural Research Center (ARC), Giza 12112, Egypt

<sup>2</sup> Benha University, Faculty of Agriculture, Soils and Water Department

Corresponding e-mail: Mohamed.abbas@fagr.bu.edu.eg (MHH Abbas) , kamelgihan @ gmail.com (G Kamel)



### **ABSTRACT**

Nine hundred thousand hectares of the irrigated lands in Egypt are salt affected soils. Such conditions threaten not only the sustainability of land use for crops production, but also the whole ecosystem. The current study aimed at investigating the growth performance of some important crops in Egypt cultivated in saline sodic clayey soils of El-Hossainia and El-Fayoum soils. To attain the aim of the study, field experiments were conducted in the investigated soils during the summer and winter seasons of the years 2013 and 2014. Rice followed by sugar beet were grown in the saline sodic soil of El-Hossainia soil; whereas sunflower followed by wheat were grown in the saline sodic soil of El-Fayoum. The amendments involved gypsum at a rate of 100% of the gypsum requirements (GR), compost at its recommended dose (100%) for each soil, a combination between gypsum at 50% of the GR and compost at 50% of its recommended dose, beside of diluted H<sub>2</sub>SO<sub>4</sub> at a rate equivalent to GR and finally bio-treatment with halophytic bacteria (Biotoul). Selected physical and chemical properties i.e. soil-water retention, soil bulk density, soil hydraulic conductivity, organic matter, soil pH and EC were considered in this study. Crop yield and growth performance of the grown plants were determined at the end of the growing seasons. The sole application of gypsum or compost at their recommended rates as well as the combined application of these amendments at 50% of these rates improved physical and chemical properties of the investigated saline sodic soils. Consequently, these treatments resulted in higher growth and growth parameters as compared with the other treatments i.e. diluted sulphuric acid and Biotoul.

**Keywords :** Saline sodic soil, soil amendments, rice, sugar beet, sun flower, wheat

### **INTRODUCTION**

Nine hundred thousand hectares of the irrigated lands in Egypt are salt affected soils (Abou-Baker and El-Dardiry, 2016). Such conditions threaten not only the sustainability of land use for crops production (Dionisio-Sese and Tobita, 1998; Pitman and Läuchli, 2002), but also the whole ecosystem (Kotb *et al.*, 2000). There are three types of salt-affected soils: (1) saline soils (EC > 4 dSm<sup>-1</sup>) (2) sodic soils (exchangeable sodium percentage (ESP) > 15) and (3) saline-sodic (EC > 4 dSm<sup>-1</sup> and ESP > 15) (Eynard *et al.*, 2005). To improve the crop yield for plants grown in saline soils, leaching is thought to be the most effective method for removing excess salts from the root zone (Abrol *et al.*, 1988). Egypt is moving towards water scarcity after the construction of the Grand Ethiopian Renaissance Dam (Mustafa *et al.*, 2013) and farmers, especially those at the end of the irrigation canals, suffer from lack of fresh water available for irrigation rather than finding sufficient water for the leaching process (Farid *et al.*, 2014). Under such conditions, saline soils could be reclaimed using biological or mechanical techniques (Ravindran *et al.*, 2007). It is thought that the agricultural soils of Egypt are going towards salinity and Sodicy. The only way for sustaining crop production in Egypt is to select salt tolerant cultivars and recommend appropriate amendments to lessen, to some extent, the hazardous effects of the soil salinity and sodicy on the grown plants.

Sodic soils contain excess sodium (Na<sup>+</sup>) that negatively affect soil structure and crop yield productivity (Qadir *et al.*, 2001). Some amendments were found to be effective in ameliorating sodic soils. Gypsum (Ca<sub>2</sub>SO<sub>4</sub>.2H<sub>2</sub>O) is a low cost material that can be used effectively in reclamation of sodic soil (Amezketta *et al.*, 2005; Makoi and Verplancke, 2010) to replace exchangeable Na in soil (Guo *et al.*, 2006). Another

effective amendment is organic matter. Organic matter can dissolve insoluble calcium salts in soils (Yamada *et al.*, 2003) to replace the exchangeable sodium ions, which are then leached out of the root zone (Ilyas *et al.*, 1997) beside of neutralizing the residual sodium carbonate in soil, to reduce pH (Choudhary *et al.*, 2011). Sulfuric acid is a well-known amendment that can prevent soil crusting and reclaim sodic soils (Amezketta *et al.*, 2005). Microbial amelioration is a promising approach for reclaiming sodic soils (Sahin *et al.*, 2011) through selection of suitable halophytes (Rabhi *et al.*, 2008).

Vegetative bioremediation “depends on growing appropriate plant species that can tolerate ambient soil salinity and Sodicy levels (Qadir and Oster, 2004). A moderate tolerant crop e.g. maize (*Zea mays* L.) (Yin *et al.*, 2004) was used effectively in soil reclamation when applying appropriate techniques (Makoi and Ndakidemi, 2007). These techniques include using soil amendments (manures and gypsum, etc.), and selecting salt-tolerant cultivars (Eynard *et al.*, 2005). Probably, plants can increase their tolerance towards salinity through increasing the rate of formation of soil aggregate and thus improve soil porosity. It is well known that plant roots can “enmesh particles together while realigning them and releasing organic compounds that hold particles together (Bronick and Lal, 2005) beside increase the stability of soil aggregates (Six *et al.*, 2000). The current study aimed at investigating the performance of the salt-tolerant important crops in Egypt grown in sodic clayey soils amended with different soil amendments.

### **MATERIALS AND METHODS**

#### **Materials of study**

##### **1-Soils**

Surface soil samples (0-30 cm) were collected from south El-Hosainiya plain (north east of the Delta region, Sharqueya Governorate), and Tamia experimental station (Tamia, El-Fayoum governorate) to represent saline sodic

soils of different physical and chemical properties. Samples were air dried, crushed, sieved to pass through a 2.0mm sieve and analyzed for their physical and chemical

properties according to the standard methods outlined by Page et al.(1982) and Klute (1986). Physical and chemical properties of the investigated soils are recorded in Table1.

**Table 1. Physical and chemical characteristics of the investigated soils**

Character	EC, dS m <sup>-1</sup>	pH	SP %	OM, g kg <sup>-1</sup>	CO <sub>3</sub> g kg <sup>-1</sup>	Particle size distribution percent					
						ESP	Clay	Silt	Fine Sand	Coarse Sand	Textural class
El – Hossainia	14.12	8.15	90	2.15	21.5	19.29	65.52	25.33	6.23	2.92	Clay
El-Fayoum	8.40	8.11	70	2.10	68.4	27.54	41.35	38.15	14.15	6.35	Clay loam

## 2-Compost

Compost was brought from El- Khalil Company, El-Khatatba under a trade name of El-Khalil compost. Its chemical and physical properties are presented in Table 2. Two rates of compost were used in this study i.e. 36 m<sup>3</sup> ha<sup>-1</sup> corresponding to the recommended dose applied in the area of study and 18 m<sup>3</sup> ha<sup>-1</sup> corresponding to 50% of the recommended dose of compost.

**Table 2. Some characteristics of El-Khalil Compost**

Property	Value
EC	dS m <sup>-1</sup> (1:10) 3.38
pH	(1:10) 7.55
Bulk density	Mg m <sup>-3</sup> 0.584
Organic matter	g kg <sup>-1</sup> 330.0
Organic Carbon	g kg <sup>-1</sup> 207.9
C/N ratio	ratio 21.43
Total N	(%) 0.97
Total P	(%) 1.23
Total K	(%) 1.34
DTPA-extractable Fe	(mg/kg) 2100.00
DTPA- extractable M n	(mg/kg) 257.00
DTPA- extractable Zn	(mg/kg) 59.00
DTPA- extractable Cu	(mg/kg) 25.00

## 3-Gypsum

Gypsum(purity 85%) was obtained from Army Chemical Factory (Koam Oshem branch, El-Fayoum). Gypsum requirements were calculated using Schoonover method(1952). The estimated values were 46 Mg gypsum ha<sup>-1</sup> recommended for the reclamation of El-Fayoum soil and 28 Mg ha<sup>-1</sup> recommended for the reclamation of El-Hosania soil. Gypsum was then broadcasted on the surface soil and then mixed thoroughly with the uppermost surface soil layer (30 cm).

## 4-Sulphuric acid

Sulphuric acid (density 1.84 Mg m<sup>-3</sup>, MW= 98.09, CAS Registry Number: 7664-93-9) was used in the current study. Sulfuric acid was applied to the investigated soils diluted with irrigation water at rates equivalent to the gypsum requirements in the investigated soils.

## 5- Biotoul compound

Biotoul compound, obtained from Soil, Water and Environmental Research Institute (SWRI), ARC, Giza, Egypt consisted of 0.98% N, 0.45% P, 0.96% Cu, 0.07% Fe, 0.39% Mn, 0.17 % Zn and some bacterial strains i.e. Azotobacter chroococcum, Azospirillum, lipoferrum, Phosphate dissolving Bacteria (Bacillus megaterium). Seeds of crops of the bio-treatments were inoculated with Biotoul compound.

## Field experiment

### El-Hossainia field experiment

Seeds of rice (*Oryza sativa* L. Giza 178) were grown during the summer season of year 2013. Plants received NPK fertilizers at recommended doses of the Ministry of Agriculture i.e. 167kg N ha<sup>-1</sup> as (NH<sub>4</sub>)<sub>2</sub>SO

(20.5% N), 16 kg P ha<sup>-1</sup> as calcium super phosphate (6.6% P) and 48 kg K ha<sup>-1</sup> as potassium sulfate (40% K). Sugar beet (*Beta vulgaris*, Diema-Fernsh) was grown in the investigated soil during the winter season of 2013/2014. NPK fertilizers were applied to soil at rates of 167 kg N ha<sup>-1</sup> as (NH<sub>4</sub>)<sub>2</sub>SO, 16 kg P ha<sup>-1</sup> as calcium super phosphate and 96kg K ha<sup>-1</sup> respectively as recommended by the Egyptian Ministry of Agriculture. All the agricultural recommended practices were followed as usual.

### El-Fayoum soil

Sun Flower (*Helianthus annuus*, Giza 102) seeds were grown during the summer season of year 2013. NPK fertilizers were applied at the recommended rates of the Ministry of Agriculture i.e. 107 kg N ha<sup>-1</sup>, 35 kg P ha<sup>-1</sup> and 48 kg K ha<sup>-1</sup>.

Wheat (*Triticum vulgare*, Sakha 94) was grown during the winter season of years 2013/2014. NPK were applied at rates of 214 kg N ha<sup>-1</sup>, 16 kg P ha<sup>-1</sup> and 40 kg K ha<sup>-1</sup>. All the agricultural practices were followed as usual recommended.

## Plant analysis

At the physiological maturity growth stage, plants were harvested, the parameters of the plant growth and crop yield for each treatment were determined. Oil in sunflower seeds was determined by Soxhlet apparatus after being extracted with petroleum ether (A.O.A.C., 1975). Juice of sugar beet was extracted from fresh macerated roots using lead acetate and its content of sucrose was determined polarimetrically (Mustafa et al., 2013). Carbohydrates % in beet roots were determined according to the Official Analysis Method (A.O.A.C., 2005).

## Data analysis

Data were statistically analyzed using the SPSS statistical software program through analysis of variance (ANOVA) and Duncan test at 0.05 probability level. For the grain crops i.e. rice and wheat crops, harvest index (HI) was calculated as the ratio of harvested grain to total shoot dry matter (Unkovich *et al.*, 2010).

# RESULTS

## Soil-water retention as affected by the different soil amendments and plants grown thereon

Results shown in Table 3 reveal that ameliorating the investigated soils with either compost, gypsum, diluted H<sub>2</sub>SO<sub>4</sub> or Biotoul improved significantly soil-water retention during both the summer and winter seasons. The highest soil-water retention values were found in soils treated with gypsum; whereas, the least ones were detected in soils treated with Biotoul. Application of “50% compost + 50% gypsum” resulted in significantly lower values of soil-water retention than

those attained with amelioration with either compost or gypsum solely. It is worthy to mention that the effect of the grown plant during reclamation seemed to be

negligible and that the amelioration effect is mainly related to the types and rates of the used treatments.

**Table 3. Soil moisture contents (%) as affected by the different soil amendments and plants grown thereon.**

	Soil tension (Bar)													
	0.001	0.01	0.33	0.66	1	3	15	0.001	0.01	0.33	0.66	1	3	15
	El-Hosania soil													
	Rice (Summer season)							Sugar beet (Winter season)						
100% CMPT	68.50	65.30	33.80	39.20	37.20	31.30	18.90	68.80	65.75	32.80	39.00	33.30	31.00	18.30
100% GPSM	66.60	63.70	34.40	38.60	33.70	30.90	18.30	66.70	63.80	34.00	38.20	36.80	30.50	18.00
50% CMPT+ 50% GPSM	64.30	62.20	33.20	37.20	33.20	29.60	19.10	64.95	62.50	32.80	36.80	32.90	29.20	19.00
H <sub>2</sub> SO <sub>4</sub>	62.80	60.70	31.70	36.50	31.90	28.90	19.90	62.90	60.90	31.50	36.20	31.50	28.50	19.80
Biotoul	61.60	59.80	31.10	36.30	31.60	28.60	20.00	61.70	59.90	31.20	36.00	31.20	28.30	20.00
Control	58.60	56.80	29.60	36.10	31.50	28.50	20.00	57.30	55.60	26.90	36.00	31.20	28.00	17.50
	El-Fayoum soil													
	Sun flower (Summer season)							Wheat (Winter season)						
100% CMPT	57.80	30.20	28.60	28.00	24.60	22.20	18.00	56.70	29.40	27.80	29.10	27.30	23.60	17.80
100% GPSM	62.70	34.70	31.90	31.70	26.80	24.70	18.70	64.60	35.30	32.80	29.90	28.20	24.00	19.80
50% CMPT+ 50% GPSM	68.60	37.60	34.20	33.10	29.40	27.10	19.80	69.10	37.90	35.00	30.60	29.30	24.40	21.00
H <sub>2</sub> SO <sub>4</sub>	63.50	34.10	31.50	29.80	26.10	24.50	18.70	64.20	34.40	31.80	29.60	28.00	23.80	19.60
Biotoul	60.80	32.90	30.90	29.60	24.90	23.40	18.40	61.70	33.40	31.40	29.10	27.10	23.40	19.40
Control	58.40	30.60	28.80	29.30	24.60	22.40	18.00	59.80	32.00	30.00	29.20	27.40	23.60	19.00

CMPT: compost; GPSM: gypsum

**Soil hydraulic conductivity as affected by the different soil amendments and plants grown thereon**

Table 4 shows that the application of compost or gypsum solely to the investigated soils increased significantly soil hydraulic conductivity. The increases occurred with application of either 100% compost or 100% gypsum were significantly higher than those

attained with application of “50% compost + 50% gypsum”. It is worthy to find that the hydraulic conductivity, in general, was higher during the winter season than the corresponding values during the summer ones. Also, Biotoul treatment increased significantly the hydraulic conductivity of the investigated soils.

**Table 4. Soil hydraulic conductivity (cm h<sup>-1</sup>) as affected by the different soil amendments and plants grown thereon**

Treatment	EL – Hossainia		EL-Fayoum	
	Rice	Sugar Beet	Sun Flower	Wheat
100% compost (CMPT)	0.086 <sup>a</sup>	0.12 <sup>b</sup>	0.06 <sup>c</sup>	0.11 <sup>b</sup>
100% Gypsum (GPSM)	0.076 <sup>b</sup>	0.18 <sup>a</sup>	0.08 <sup>a</sup>	0.17 <sup>a</sup>
50% CMPT + 50% GPSM	0.046 <sup>d</sup>	0.08 <sup>c</sup>	0.02 <sup>f</sup>	0.09 <sup>c</sup>
H <sub>2</sub> SO <sub>4</sub>	0.056 <sup>c</sup>	0.07 <sup>d</sup>	0.07 <sup>b</sup>	0.08 <sup>d</sup>
Biotoul	0.016 <sup>e</sup>	0.06 <sup>e</sup>	0.04 <sup>d</sup>	0.06 <sup>e</sup>
Control	0.001 <sup>f</sup>	0.03 <sup>f</sup>	0.03 <sup>e</sup>	0.05 <sup>f</sup>

**Soil bulk density as affected by the different soil amendments and plants grown there on.**

Table 5 shows that values of soil bulk density increased in the saline sodic soils with application of all treatments except for compost; however, the combined

application of “50% compost and 50% gypsum” resulted in the highest values of soil bulk density. The grown plants seemed to be of an obvious effect on values of bulk density in both the investigated soils.

**Table 5. Soil bulk density (Mg m<sup>-3</sup>) as affected by the different soil amendments and plants grown thereon**

Treatment	EL – Hossainia		EL-Fayoum	
	Rice	Sugar Beet	Sun Flower	Wheat
100% compost (CMPT)	1.21 <sup>f</sup>	1.17 <sup>f</sup>	1.20 <sup>f</sup>	1.18 <sup>f</sup>
100% Gypsum (GPSM)	1.34 <sup>b</sup>	1.24 <sup>b</sup>	1.24 <sup>c</sup>	1.22 <sup>c</sup>
50% CMPT + 50% GPSM	1.41 <sup>a</sup>	1.30 <sup>a</sup>	1.33 <sup>a</sup>	1.28 <sup>a</sup>
H <sub>2</sub> SO <sub>4</sub>	1.30 <sup>c</sup>	1.23 <sup>c</sup>	1.29 <sup>b</sup>	1.24 <sup>b</sup>
Biotoul	1.26 <sup>d</sup>	1.21 <sup>d</sup>	1.22 <sup>d</sup>	1.20 <sup>d</sup>
Control	1.24 <sup>e</sup>	1.18 <sup>e</sup>	1.21 <sup>e</sup>	1.19 <sup>e</sup>

**Soil organic matter as affected by the different soil amendments and plants grown thereon.**

Application of compost or gypsum solely or together at 50% of their recommended rates to the studied soils resulted in relatively higher contents of organic matter in soils compared with the other treatments (Table 6). H<sub>2</sub>SO<sub>4</sub> and Biotoul treatments resulted in the lowest significant increases in soil

organic matter. The soil organic matter also varied according to the grown plant type.

**EC and pH of the soils treated with different soil amendments.**

Table 7 shows that application of compost solely or H<sub>2</sub>SO<sub>4</sub> to the studied soils decreased significantly soil pH and EC. Although, the application of gypsum decreased soil EC and pH in Hossainia soil; however, in

Fayoum soil, the results seemed confusing especially that the application of gypsum decreased slightly soil pH, with no significant effect on soil EC. Biotoul was found to be of the least effect on both soil pH and EC in

El-Hossainia soil, yet, it was ineffective in minimizing such parameters in El-Fayoum soil.

**Table 6. Soil organic matter (g kg<sup>-1</sup>) as affected by the different soil amendments and plants grown thereon.**

Treatment	EL – Hossainia		EL-Fayoum	
	Rice	SugarBeat	SunFlower	Wheat
100% compost (CMPT)	4.7 <sup>a</sup>	4.6 <sup>a</sup>	4.4 <sup>a</sup>	4.2 <sup>a</sup>
100% gypsum (GPSM)	3.5 <sup>c</sup>	3.4 <sup>c</sup>	3.3 <sup>c</sup>	3.3 <sup>c</sup>
50% CMPT + 50% GPSM	4.2 <sup>b</sup>	4.2 <sup>b</sup>	4.3 <sup>b</sup>	4.1 <sup>b</sup>
H <sub>2</sub> SO <sub>4</sub>	2.7 <sup>d</sup>	2.5 <sup>d</sup>	2.5 <sup>d</sup>	2.2 <sup>d</sup>
Biotoul	2.3 <sup>e</sup>	2.2 <sup>e</sup>	2.4 <sup>e</sup>	2.2 <sup>d</sup>
Control	2.1 <sup>f</sup>	2.1 <sup>f</sup>	2.4 <sup>c</sup>	2.2 <sup>d</sup>

**Table 7. EC and pH of the investigated soils treated with different soil amendments**

Treatment	El-Hossainia		El-Fayoum	
	EC <sub>e</sub> dS m <sup>-1</sup>	pH	EC <sub>e</sub> dS m <sup>-1</sup>	pH
100% compost (CMPT)	7.27 <sup>d</sup>	7.66 <sup>cd</sup>	7.46 <sup>b</sup>	7.78 <sup>c</sup>
100% gypsum (GPSM)	6.26 <sup>f</sup>	7.70 <sup>bcd</sup>	8.36 <sup>a</sup>	7.87 <sup>b</sup>
50% CMPT + 50% GPSM	9.86 <sup>c</sup>	7.61 <sup>d</sup>	7.17 <sup>c</sup>	7.71 <sup>cd</sup>
H <sub>2</sub> SO <sub>4</sub>	6.58 <sup>e</sup>	7.70 <sup>bcd</sup>	6.86 <sup>d</sup>	6.70 <sup>e</sup>
Biotoul	9.82 <sup>c</sup>	7.78 <sup>b</sup>	8.37 <sup>a</sup>	7.94 <sup>ab</sup>
Control	14.11 <sup>a</sup>	8.13 <sup>a</sup>	8.39 <sup>a</sup>	8.01 <sup>a</sup>

**Growth parameters and yield components of plants grown in saline-sodic soils treated with different soil amendments**

Application of compost to the investigated soils recorded the highest significant increases in the growth and growth parameters of both rice and sun flower (Table 8). On the other hand, application of gypsum to the investigated soils recorded the highest increases in the growth parameters of sugar Beet plants beside of its superior effects on the grain and straw yields of the winter wheat. Treating soils with either H<sub>2</sub>SO<sub>4</sub> or

Biotoul recorded slight; however, significant increases in the growth parameters of all crops under study except for rice.

Carbohydrates % of sugar beet and oil % of sun flower slightly increased due to all the studied treatments except for Biotoul; however, such increases were significant. Values of harvest index of both rice and wheat slightly flocculated between 0.57- 0.64 and 1.33-2.30, respectively. As a matter of fact, these ranges of flocculation seemed somewhat lower than the corresponding ones of the control treatments.

**Table 8. Growth parameters and yield components of plants grown in the saline-sodic soils treated with different soil amendments**

Treatment	Rice				Sugar beet			
	Grain yield Mg.ha <sup>-1</sup>	Straw yield, Mg ha <sup>-1</sup>	100-grain weight	Harvest Index	Weight Mg.ha <sup>-1</sup>	Diameter of sugar beet, cm	Carbohyd rate, %	Sucrose %
100% CMPT	7.05 <sup>a</sup>	11.24 <sup>a</sup>	1.49 <sup>a</sup>	0.63	43.57 <sup>c</sup>	19.93 <sup>a</sup>	0.12 <sup>a</sup>	21 <sup>a</sup>
100% GPSM	6.81 <sup>b</sup>	10.83 <sup>b</sup>	1.67 <sup>b</sup>	0.63	45.00 <sup>a</sup>	18.98 <sup>b</sup>	0.12 <sup>a</sup>	20 <sup>b</sup>
50%CMPT+ 50%GPSM	6.33 <sup>c</sup>	9.91 <sup>c</sup>	1.45 <sup>c</sup>	0.64	40.95 <sup>b</sup>	17.99 <sup>c</sup>	0.11 <sup>b</sup>	19 <sup>c</sup>
H <sub>2</sub> SO <sub>4</sub>	2.76 <sup>d</sup>	4.83 <sup>d</sup>	1.09 <sup>d</sup>	0.57	40.95 <sup>d</sup>	15.99 <sup>d</sup>	0.11 <sup>b</sup>	17 <sup>d</sup>
Biotoul	2.76 <sup>d</sup>	4.52 <sup>f</sup>	1.06 <sup>d</sup>	0.61	40.24 <sup>e</sup>	14.99 <sup>e</sup>	0.11 <sup>b</sup>	16 <sup>e</sup>
Control	2.76 <sup>d</sup>	4.19 <sup>f</sup>	1.06 <sup>d</sup>	0.66	40.23 <sup>c</sup>	11.96 <sup>f</sup>	0.10 <sup>c</sup>	15 <sup>f</sup>

  

Treatment	Sun flower				Wheat			
	Yield	Length of Stalk, cm	Diameter of disk, Cm	Oil, %	grain Mg.ha <sup>-1</sup>	straw Mg.ha <sup>-1</sup>	100-grain weight	Harvest Index
100% CMPT	60.45 <sup>a</sup>	114.66 <sup>c</sup>	32.66 <sup>b</sup>	60.45 <sup>a</sup>	3.16 <sup>a</sup>	2.30 <sup>a</sup>	1.92 <sup>a</sup>	1.37
100% GPSM	55.12 <sup>c</sup>	119.66 <sup>b</sup>	34.66 <sup>a</sup>	55.12 <sup>c</sup>	3.33 <sup>b</sup>	2.50 <sup>f</sup>	1.95 <sup>b</sup>	1.33
50%CMPT+ 50%GPSM	55.52 <sup>b</sup>	97.66 <sup>d</sup>	31.66 <sup>c</sup>	55.52 <sup>b</sup>	2.50 <sup>c</sup>	1.50 <sup>b</sup>	1.79 <sup>c</sup>	1.67
H <sub>2</sub> SO <sub>4</sub>	51.72 <sup>d</sup>	73.66 <sup>e</sup>	29.66 <sup>d</sup>	51.72 <sup>d</sup>	2.45 <sup>d</sup>	1.20 <sup>c</sup>	1.43 <sup>d</sup>	2.04
Biotoul	51.36 <sup>e</sup>	65.66 <sup>f</sup>	26.66 <sup>e</sup>	51.36 <sup>e</sup>	2.30 <sup>e</sup>	1.00 <sup>d</sup>	1.40 <sup>e</sup>	2.30
Control	51.36 <sup>e</sup>	159.66 <sup>a</sup>	24.66 <sup>f</sup>	51.36 <sup>e</sup>	2.30 <sup>f</sup>	0.98 <sup>e</sup>	1.25 <sup>f</sup>	2.35

CMPT: compost; GPSM: gypsum

**DISCUSSION**

Sodic soil is characterized by the presence of high concentrations of sodium (Na) as an exchangeable cation (Sumner, 1993) causing clay dispersion (Campos et al., 2006; Qadir et al., 2007), deterioration of soil hydraulic properties (Qadir and Schubert, 2002) and adversely affect the plant growth (Dagar et al., 2001; Akhter et al., 2003; Qadir et al., 2003). Thus,

ameliorating such soils is the only mean to improve soil properties and attain better crop yield. Plant assisted remediation is a promising approach to replace the more costly chemical approaches (Qadir et al., 2002). Soil organic carbon (SOC) pool should also be considered in soil reclamation studies because SOC is an important measure of soil health (Wong et al., 2010). Hydraulic conductivity is one of the important parameters to be considered in the hydrological cycle (Candemir and

Gülser, 2012). Application of compost or gypsum to the studied soils improved soil-water retention and, at the same time, improved soil hydraulic conductivity. Generally, the residual soil organic matter increased but the soil bulk density increased in such soils. However, such changes in EC and bulk density differed, to some extent, from a crop to another. Accordingly, it can be deduced that all growth parameters and yield components of the cultivated crops improved with applying such amendments.

#### **Compost for ameliorating soil salinity and sodicity**

Results indicate that application of the organic matter improved soil physical properties, decreased soil EC and pH and increased plant growth and yield components in the investigated soils, especially at its highest rates i.e. 100% OM. Compost applications released acids which could ultimately increase the solubility of soil CaCO<sub>3</sub> to replace the exchangeable sodium by calcium (Avnimelech *et al.*, 1994). Thus, reduction in pH and ESP values occurred due to application of the compost to the alkali (sodic) soils (Rao and Pathak, 1996). Also, organic matter increases the formation of soil aggregates (Six *et al.*, 2004; Abbas *et al.*, 2012) and stabilizes soil structure and enhances plant growth (Avnimelech *et al.*, 1994; Puget *et al.*, 2000) beside of stimulating soil microbial biomass (SMB) (Wong *et al.*, 2009).

#### **Gypsum for ameliorating soil salinity and sodicity**

Gypsum applications improved soil physical properties and increased the crop yield in investigated sodic soils. Gypsum supplied calcium (Ca<sup>2+</sup>) directly to soils to replace excess exchangeable sodium (Na<sup>+</sup>) (Qadir and Oster, 2002) and; therefore, improved the infiltration rate and hydraulic conductivity in soils (Sahin *et al.*, 2003; Gharaibeh *et al.*, 2009; Reading *et al.*, 2012; Batool *et al.*, 2015). Accordingly, leaching of salts from soils was improved (Gharaibeh *et al.*, 2009). It is worthy to mention that the internal soil swelling increased water holding capacity of the soil at low electrolyte concentrations (Mace and Amrhein, 2001). Such conditions could improve the crop growth (Hamza and Anderson, 2002; Yao *et al.*, 2013), encouraging further root exudates to increase formation of soil aggregations (Angers and Caron, 1998), in addition to the residual organic carbon in soils resulted from increasing crop yield. Thus, gypsum increases water retention in soils (Moret-Fernández and Herrero, 2015).

No reductions occurred in soil pH and EC with gypsum applications to El-Fayoum soil. It seems that the mode of action of gypsum in El-Fayoum soil is, mainly due to antagonizing the effect of Na and reducing its uptake by plants (Navarro *et al.*, 2000; Montesano and van Iersel, 2007). Accordingly, the plant growth parameters and yield components of sun flower and wheat plants grown in El-Fayoum soil were lower than those attained due to application of compost to the soil.

#### **Application of diluted H<sub>2</sub>SO<sub>4</sub> for ameliorating soil salinity and sodicity**

Although soil pH and EC decreased with applying diluted sulphuric acid with irrigation water, and that the measured values of pH and EC were lower than

those attained due to applications of either gypsum or compost; however, the corresponding increases in the crop yield components which were treated with the acidified irrigation water were significantly lower than those attained with applications of either gypsum or compost. Such a result probably indicates that the ameliorating effect of the compost exceeded the effect of just acidifying soil rhizosphere which increased the solubility of insoluble CaCO<sub>3</sub> found in the soil. Diluted acids can solubilize the native CaCO<sub>3</sub> to supply sufficient Ca<sup>2+</sup> for reclamation of sodic soils (Qadir *et al.*, 1996). Probably the grown plants played significant roles in the process of ameliorating the investigated saline sodic soils.

#### **Biological approach to improve crop yield grown in saline- sodic soils**

Slight or insignificant changes took place in the abovementioned physical and chemical characteristics of the investigated soils with application of Biotoul. The crop performance slightly changed with Biotoul application. Such a result probably indicates that increasing plant tolerance towards salinity sodicity is not enough to attain better crop yield without further improvements in soil physical and chemical characteristics.

## **REFERENCES**

- Abbas, M.H.H., Ismael, A.O.A., El-Gamal, M.A.H., Salem, H.M. 2011. Integrated effect of mineral nitrogen, bio and organic fertilization on soybean productivity Egypt J Biotechnol 39, 43-63
- Abbas, G., Saqib, M., Rafique, Q., ur-Rahman, M.A., Akhtar, J., Abdul-Haq, M. and Nasim, M. 2013. Effect of salinity on grain yield and grain quality of wheat (*Triticum aestivum*L.). *Pak. J. Agri. Sci.*, 50: 185-189.
- Abou-Baker, N., El-Dardiry, E. 2016. Integrated management of salt affected soils in agriculture: Incorporation of soil salinity control methods. Academic Press, Elsevier, London.
- Abrol, I.P., Yadav, J.S.D., Massoud, F.I., 1988. Salt-affected soils and their management. Food and Agriculture Organization of the United Nations. Bulletin 39. Available at: <http://www.fao.org/docrep/x5871e/x5871e00.htm>
- Akhter, J., Mahmood, M., Malik, K.A., Ahmed, S., 2003. Amelioration of a saline sodic soil through cultivation of a salt-tolerant grass *Leptochloa fusca*. *Environmental Conservation*, 168-174.
- Amezketta, E., Aragüés, R., Gazol, R., 2005. Efficiency of sulfuric acid, mined gypsum, and two gypsum by-products in soil crusting prevention and sodic soil reclamation. *Agronomy Journal* 97, 983-989.
- Angers, D.A., Caron, J., 1998. Plant-induced changes in soil structure: Processes and feedbacks. In: van Breemen, N. (Ed.), *Plant-induced soil changes: Processes and feedbacks*. Springer-Science+Business Media, B.V., pp. 55-72.
- AOAC (1975): *Official Methods of Analysis*, 12th ed., Association of Official Analytical Chemists, Washington, DC.

- AOAC (2005). Official Methods of Analysis. Association of Official Analytical Chemists, Washington DC.
- Amezqueta, E., Aragués, R. and Gazol, R., 2005. Efficiency of sulfuric acid, mined gypsum, and two gypsum by-products in soil crusting prevention and sodic soil reclamation *Agronomy Journal* 97, 983-989.
- Avnimelech, Y., Shkedy, D., Kochva, M., Yotal, Y., 1994. The Use of Compost for the Reclamation of Saline and Alkaline Soils. *Compost Science & Utilization* 2, 6-11.
- Batool, A., Taj, S., Rashid, A., Khalid, A., Qadeer, S., Saleem, A.R., Ghufuran, M.A., 2015. Potential of soil amendments (Biochar and Gypsum) in increasing water use efficiency of *Abelmoschus esculentus* L. Moench. *Frontiers in Plant Science* 6, 733.
- Bronick, C.J., Lal, R., 2005. Soil structure and management: a review. *Geoderma* 124, 3-22.
- Campos, C.A.B., Fernandes, P.D., Gheyi, H.R., Blanco, F.F., Gonçalves, C.B., Campos, S.A.F., 2006. Yield and fruit quality of industrial tomato under saline irrigation. *Scientia Agricola* 63, 146-152.
- Candemir, F., Gülser, C., 2012. Influencing factors and prediction of hydraulic conductivity in fine-textured alkaline soils. *Arid Land Research and Management* 26, 15-31.
- Choudhary, O.P., Ghuman, B.S., Bijay, S., Thuy, N., Buresh, R.J., 2011. Effects of long-term use of sodic water irrigation, amendments and crop residues on soil properties and crop yields in rice-wheat cropping system in a calcareous soil. *Field Crops Research* 121, 363-372.
- Dagar, J.C., Singh, G., Singh, N.T., 2001. Evaluation of forest and fruit trees used for rehabilitation of semiarid alkali-sodic soils in India. *Arid Land Research and Management* 15, 115-133.
- Dionisio-Sese, M.L., Tobita, S., 1998. Antioxidant responses of rice seedlings to salinity stress. *Plant Science* 135, 1-9.
- Eynard, A., Lal, R., Wiebe, K., 2005. Crop response in salt-affected soils. *Journal of Sustainable Agriculture* 27, 5-50.
- Farid, I.M., Abbas, M.H.H., Fawzy, E., 2014. Rationalizing the use of water of salinity hazards for irrigating maize grown in a saline sodic soil. *Egypt. J. Soil Sci.* 54, 163-175.
- Gharaibeh, M.A., Eltaif, N.I., Shunnar, O.F., 2009. Leaching and reclamation of calcareous saline-sodic soil by moderately saline and moderate-SAR water using gypsum and calcium chloride. *Journal of Plant Nutrition and Soil Science* 172, 713-719.
- Guo, G., Araya, K., Jia, H., Zhang, Z., Ohomiya, K., Matsuda, J., 2006. Improvement of salt-affected soils, Part 1: Interception of capillarity. *Biosystems Engineering* 94, 139-150.
- Hamza, M.A., Anderson, W.K., 2002. Improving soil physical fertility and crop yield on a clay soil in Western Australia. *Australian Journal of Agricultural Research* 53, 615-620.
- Ilyas, M., Qureshi, R.H., Qadir, M.A., 1997. Chemical changes in a saline-sodic soil after gypsum application and cropping. *Soil Technology* 10, 247-260.
- Klute, A., 1986. Part 1. Physical and mineralogical methods. ASA-SSSA-Agronomy, Madison, Wisconsin USA.
- Kotb, T.H.S., Watanabe, T., Ogino, Y., Tanji, K.K., 2000. Soil salinization in the Nile Delta and related policy issues in Egypt. *Agricultural Water Management* 43, 239-261.
- Mace, J.E., Amrhein, C., 2001. Leaching and reclamation of a soil irrigated with moderate SAR waters. *Soil Science Society of America Journal* 65.
- Makoi, J., Ndakidemi, P., 2007. Reclamation of sodic soils in northern Tanzania, using locally available organic and inorganic resources. *African Journal of Biotechnology* 6, 1926-1931.
- Makoi, J.H., Verplancke, H., 2010. Effect of gypsum placement on the physical chemical properties of a saline sandy loam soil. *Australian Journal of Crop Science* 4, 556-563.
- Montesano, F., van Iersel, M.W., 2007. Calcium can prevent toxic effects of Na<sup>+</sup> on tomato leaf photosynthesis but does not restore growth. *Journal of the American Society for Horticultural Science* 132, 310-318.
- Moret-Fernández, D., Herrero, J., 2015. Effect of gypsum content on soil water retention. *Journal of Hydrology* 528, 122-126.
- Mustafa, E.F., Farid, I.M., Abbas, M.H.H., 2013. Yield economical return and amelioration effect of sugar beet grown in sodic soil irrigated with low quality water 2.
- Navarro, J.M., Botella, M.Á., Cerdá, A., Martínez, V., 2000. Effect of salinity × calcium interaction on cation balance in melon plants grown under two regimes of orthophosphate. *Journal of Plant Nutrition* 23, 991-1006.
- Page, A.L., Miller, R.H., Keeney, D.R., 1982. Methods of soil analysis Part II Chemical and microbiological properties ASA-SSSA. Agronomy, Madison, USA.
- Pitman, M.G., Läuchli, A., 2002. Global Impact of Salinity and Agricultural Ecosystems. In: Läuchli, A., Lüttge, U. (Eds.), *Salinity: Environment - Plants - Molecules*. Springer Netherlands, Dordrecht, pp. 3-20.
- Puget, P., Chenu, C., Balesdent, J., 2000. Dynamics of soil organic matter associated with particle-size fractions of water-stable aggregates. *European Journal of Soil Science* 51, 595-605.
- Qadir, M., Oster, J., 2002. Vegetative bioremediation of calcareous sodic soils: history, mechanisms, and evaluation. *Irrigation Science* 21, 91-101.
- Qadir, M., Oster, J.D., 2004. Crop and irrigation management strategies for saline-sodic soils and waters aimed at environmentally sustainable agriculture. *Science of The Total Environment* 323, 1-19.
- Qadir, M., Oster, J.D., Schubert, S., Noble, A.D., Sahrawat, K.L., 2007. Phytoremediation of sodic and saline - sodic soils. *Advances in Agronomy*. Academic Press, pp. 197-247.

- Qadir, M., Qureshi, R.H., Ahmad, N., 1996. Reclamation of a saline-sodic soil by gypsum and *Leptochloa fusca*. *Geoderma* 74, 207-217.
- Qadir, M., Qureshi, R.H., Ahmad, N., 2002. Amelioration of calcareous saline sodic soils through phytoremediation and chemical strategies. *Soil Use and Management* 18, 381-385.
- Qadir, M., Schubert, S., 2002. Degradation processes and nutrient constraints in sodic soils. *Land Degradation & Development* 13, 275-294.
- Qadir, M., Schubert, S., Ghafoor, A., Murtaza, G., 2001. Amelioration strategies for sodic soils: a review. *Land Degradation & Development* 12, 357-386.
- Qadir, M., Steffens, D., Yan, F., Schubert, S., 2003. Sodium removal from a calcareous saline-sodic soil through leaching and plant uptake during phytoremediation. *Land Degradation & Development* 14, 301-307.
- Rabhi, M., Talbi, O., Atia, A., Abdelly, C., Smaoui, A., 2008. Selection of a halophyte that could be used in the bioreclamation of salt-affected soils in arid and semi-arid regions. In: Abdelly, C., Öztürk, M., Ashraf, M., Grignon, C. (Eds.), *Biosaline agriculture and high Salinity tolerance*. Birkhäuser Basel, Basel, pp. 241-246.
- Rao, D.L.N., Pathak, H., 1996. Ameliorative influence of organic matter on biological activity of salt - affected soils. *Arid Soil Research and Rehabilitation* 10, 311-319.
- Ravindran, K.C., Venkatesan, K., Balakrishnan, V., Chellappan, K.P., Balasubramanian, T., 2007. Restoration of saline land by halophytes for Indian soils. *Soil Biology and Biochemistry* 39, 2661-2664.
- Reading, L.P., Baumgartl, T., Bristow, K.L., Lockington, D.A., 2012. Hydraulic conductivity increases in a sodic clay soil in response to gypsum applications: Impacts of bulk density and cation exchange. *Soil Science* 177, 165-171.
- Sahin, U., Eroğlu, S., Sahin, F., 2011. Microbial application with gypsum increases the saturated hydraulic conductivity of saline-sodic soils. *Applied Soil Ecology* 48, 247-250.
- Sahin, U., Oztas, T., Anapali, O., 2003. Effects of consecutive applications of gypsum in equal, increasing, and decreasing quantities on soil hydraulic conductivity of a saline-sodic soil. *Journal of Plant Nutrition and Soil Science* 166, 621-624.
- Six, J., Bossuyt, H., Degryze, S., Denef, K., 2004. A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research* 79, 7-31.
- Six, J., Paustian, K., Elliott, E.T., Combrink, C., 2000. Soil structure and organic matter I. Distribution of aggregate-size classes and aggregate-associated carbon. *Soil Science Society of America Journal* 64, 681-689.
- Sumner, M.E., 1993. Sodic soils - New perspectives. *Australian Journal of Soil Research* 31, 683-750.
- Unkovich, M., Baldock, J., Forbes, M., 2010. Chapter 5 - Variability in harvest index of grain crops and potential significance for carbon accounting: Examples from Australian agriculture. *Advances in Agronomy*. Academic Press, pp. 173-219.
- Wong, V.N.L., Dalal, R.C., Greene, R.S.B., 2009. Carbon dynamics of sodic and saline soils following gypsum and organic material additions: A laboratory incubation. *Applied Soil Ecology* 41, 29-40.
- Wong, V.N.L., Greene, R.S.B., Dalal, R.C., Murphy, B.W., 2010. Soil carbon dynamics in saline and sodic soils: a review. *Soil Use and Management* 26, 2-11.
- Yamada, P., N. Ohsawa, N., Yazawa, Y., Nishizawa, Y., Yamaguchi, T., 2003. Improvement of physicochemical properties of alkali soil by humic materials: addition effect of the local peat and weathered coal on the paddy field rice cultivation on the saline arid land in Keerqin desert, China. *Journal of Arid Land Studies* 12, 203-213.
- Yao, R.-J., Yang, J.-S., Zhang, T.-J., Gao, P., Yu, S.-P., Wang, X.-P., 2013. Short-term effect of cultivation and crop rotation systems on soil quality indicators in a coastal newly reclaimed farming area. *Journal of Soils and Sediments* 13, 1335.
- Yin, X.Y., Yang, A.F., Zhang, K.W., Zhang, J.R., 2004. Production and analysis of transgenic maize with improved salt tolerance by the introduction of *AtNHX1* gene. *Acta Botanica Sinica* 46, 854-861.

## التخفيف من حدة الملوحة والصودية بإضافة محسنات التربة

جيهان حسين كامل<sup>1,2</sup>، عصمت نوفل<sup>2</sup>، إيهاب محمد فريد<sup>2</sup>، سمير عبد العزيز<sup>1</sup> و محمد حسن حمزة عباس<sup>2</sup>

1- معهد بحوث الأراضي والمياه والبيئة، المركز البحوث الزراعية، الجيزة

2- قسم الأراضي والمياه، كلية الزراعة، جامعة بنها

تبلغ مساحة الارض المروية في مصر المتأثرة بالأملاح تسعة مائة الف هكتار ومثل هذه الظروف تهدد بعدم إستدامة إنتاج المحاصيل في تلك الاضي، كما تهدد النظام البيئي بأكمله، وبالتالي، تهدف الدراسة الحالية إلى التعرف على تأثير بعض محسنات التربة على الحد من ملوحة وصودية التربة وكذلك على بعض المحاصيل المنزرعة في أراضي ملحية صودية في مناطق الحسينية والفيوم، وللتحقق من هدف الدراسة، فإنه تم اجراء تجربة حقلية لمعرفة أثر هذه المحسنات على الاراضي محل الدراسة خلال فصلي الصيف والشتاء (2013- 2014) حيث تم زراعة محصول الأرز يليه محصول بنجر السكر في منطقة الحسينية بمحافظة الشرقية، كما تمت زراعة محصول عباد الشمس يليه محصول القمح في أرض ملحية قلوية في طامية بمحافظة الفيوم، وكانت معدلات إضافة محسنات التربة هي: (1) إضافة الجبس 100% من إحتياجات الجبسية للارض، (2) إضافة الكميوست بمعدل 100% من الكمية الموصي بها، (3) إضافة مزيج من 50% الكميوست : 50% الجبس من الدفعات الموصي بها، (4) إضافة حمض الكبريتيك المخفف وذلك بما يعادل الجبس الموصي اضافته للتربة عند 100% من الإحتياجات الجبسية، (5) إضافة مركب حيوي عبارة عن بكتريا مقاومة للملوحة تحت الاسم التجاري(بيوترون)، وقد تم تقدير بعض الخصائص الفيزيائية والكيميائية للتربة وهي احتجاز التربة للرطوبة تحت قيم مختلفة من الشد الرطوبي، كما قدرت الكثافة الظاهرية و التوصيل الهيدروليكي للتربة ومحتوي التربة من المادة العضوية وحموضة التربة والتوصيل الكهربي للتربة، بالإضافة إلى التقديرات المحصولية ومؤشرات نمو المحاصيل المنزرعة في نهاية موسم نموها، حيث ادت الإضافات الفردية لكل من الكميوست بمعدل 100% من الكمية الموصي بها وكذلك الجبس بمعدل 100% من الإحتياجات الجبسية إلى تحسين الخواص الفيزيائية والكيميائية للتربة الملحية الصودية وقد أعطت اعلى محصول من كل من (الأرز-بنجر السكر - عباد الشمس - القمح)، أما بالنسبة لباقي المعاملات الاخرى وهي 50% من المعدل الموصي به لكل من الكميوست والجبس بالإضافة إلى حمض الكبريتيك المخفف والمركب الحيوي(البيوترون). فكانت ذات أثر اقل على كل من التربة والمحصول.