

EFFECTIVENESS OF APPLYING NATURAL SOIL CONDITIONERS UNDER MOISTURE REGIMES ON SANDY SOIL PROPERTIES AND CEREAL CROPS PRODUCTION

EI-Kammah, M.A.M¹; Marwa, G. M. Ali²; M.A.Aboelsoud² and S. A. Mashalli¹

1-Soils and Water Sci. Dept., Fac. of Agric., Kafrelsheikh Univ., Kafr El-Sheikh, Egypt

2-Soils, Water and Environ. Res. Inst., Sakha Agric. Res. Station, Egypt

ABSTRACT

Two experimental sites of arable sandy soil located at Abou Omera Al-Sharkeya village, Baltim district, Kafr El-Sheikh Governorate were chosen. The selected locations represent the conditions of circumstances of northern part of Nile Delta region. Site 1 (fruit field) was devoted for collecting composite surface soil sample for carrying out wheat pot experiment in plastic pots during the growing winter season period 2010/2011. Site 2 (nearby site 1) was occupied for conducting maize field experiment during growing summer season period 2011. The main objectives of this investigation were to study and evaluate the effect of natural raw minerals, soil conditioner types, their mixtures and application rates in sandy soil subjected to different irrigation deficits on the following parameters: (i) soil properties (chemical, physical, moisture constants and nutritional status after cereal crops harvesting and (ii) agronomical production of wheat and maize crops after full maturity in the studied soil under consideration.

Four types of soil conditioners [bentonite, compost, mixture of natural mineral raw materials (MNRM) and their mixtures 1:1:1 (w/w)] were applied before cultivation in two recommended application rates low (R₁) and high (R₂). Three levels of soil moisture depletion regimes were used (30, 50 and 70%) from its available water capacity. The N, P, K mineral fertilizers were added according to the recommended doses for sandy soils. Applying natural soil conditioner types, their mixtures and application rates in the studied sandy soil subjected to moisture depletion regime realized improving soil chemical, physical, water holding capacity and macro nutritional status. Soil salinity (soil reaction, electrical conductivity and ionic strength), hazard sodium parameters (soluble sodium percentage and sodium adsorption ratio), soil porosity, available water capacity and phyto-available nutrients were increased. On the other hand, hydraulic conductivity and bulk density were decreased. Generally, high application rate achieved the best values of soil properties in comparison with low application rate. Conditioner mixtures 1:1:1 treatment realized the superiority under wheat pot experiment, meanwhile, compost treatment achieved the best values under maize field experiment. Irrigation after 50% AWSMD gave moderate values of such properties between wet (30% AWSMD) and dry (70% AWSMD). Increasing grains, straw, protein, grains weight, harvest index and other yield components significantly for wheat and maize crops, as a result of adding soil conditioners compared with control. Conditioner mixtures 1:1:1 treatment realized the superiority under wheat pot experiment, meanwhile, compost treatment achieved the best values under maize field experiment. Also, high application rate was better than low application rate. Results also indicated that, from view point of water and economic, the highest values of crop yield were obtained from irrigation at 50% depletion from its available water capacity rather than 30% and 70% respectively.

Keywords: Guelph permeameter apparatus; Time Domain Reflectometer (TDR apparatus); Composting; Nutrients – phytoavailability; Sandy soils; Water irrigation management; Soil conditioners; Cutthroat flumes.

INTRODUCTION

Nowadays, the term of « sustainable agriculture » is widely used in world wide, which is keystone of the rational utilization of soils as one of our most important natural resources. It is the important aims of « sustainable agriculture » to protect and maintain of the multifunctions of soils (Varallyay, 2005). For preservation and sustainability the productivity of soil we have to take special regard to sandy soils having unfavorable properties. Sandy soil characterized by less than 18 % clay and more than 68 % sand in the first 100 cm of the soil depth are the poor soils that occur in many parts of the world (van Wambeke, 1992). There are other problems facing agriculture sector caused by, mainly, inappropriate soil, water and fertile management practices as well as rapid decreasing of agricultural land particularly in Delta soils. Therefore, we have to find rapid solutions to face these problems. Sandy soils hold little water as the large pore spaces allow water to drain freely from soil. The productivity of these soils is limited by low water holding capacities, high infiltration rates, high evaporation, low inherent fertility levels, very low organic matter content and excessive deep percolation losses. Also, the water use efficiency of the crops cultivated in such soil is low.

Tackling these problems can be achieved through applying organic amendments, natural raw minerals and soil conditioners. These materials improve the retentative capacities of these soils and allow plants to get their water requirements and phyto –available nutrients easily.

Cereal crops such as (wheat and maize) are very strategically important crops in Egypt because it's constituent and indispensable part of Egyptian food diet. Generally, there is a great gap between the consumption and production of such crops. On the other hand, it is worthnoting that, the agriculture production in Egypt is mainly depend upon irrigated agriculture. The gap between supplies and demands of water is widening with increasing global population. We are suffering from this trouble, especially when we know that we are under water poverty limit. Because of the water limitation, one of the most important targets in the agriculture sector is how to save irrigation water and increase water use efficiencies. So, new techniques and practices are needed to achieve water save. Estimating irrigation water becomes important for project planning and irrigation management. The over irrigation practiced by the farmers usually leads to low irrigation efficiency. So it is necessary to ascertain to what extent the water in the root zone can be depleted to produce high economic yield with using little water applied . Planning best irrigation regimes is very important for maintaining available irrigation water. The proper water management (irrigation scheduling) not only accurates determination of crop water requirements but also helps to know when and how much water should be applied to get high efficiency of each unit of water. Regulated deficit irrigation is one of such practices. Many studies indicated that the deficit irrigation was a successful technique in crops irrigation, Omran(2005) and Seif *et al.*(2005). The main objectives of this investigation were to study and asses the effect of natural soil conditioner types, their mixtures and application rates in sandy soils subjected to irrigation regimes on:

- (i): Soil physico-chemical properties, moisture constants and nutritional status.
(ii): Agronomical production of wheat and maize crops after full maturity.

MATERIALS AND METHODS

Two selected sites represent arable sandy soil located at Abou-Omera Al-Sharkeya village, Baltim district, Kafr El-Sheikh Governorate 31° 34 40.6 N latitude and 31° 10 55.5 E longitude with an elevation of about 5 meters above sea level were chosen. Site 1 (fruit field) was devoted for collecting composite surface soil sample for carrying out wheat pot experiment in plastic pots during the growing winter season period 2010/2011. Site 2 (nearby site 1) was occupied for conducting maize field experiment during growing summer season period 2011. After wheat and maize plants full maturity, representative composite disturbed soil surface samples were collected, air dried, crumbled by hand, homogenized and finely ground in steel mill to pass through 100-mesh (0.15 mm opening sieve) and thoroughly mixed.

Generally, soil chemical characterizations of the studied soils before cultivation and directly after cereal crops harvesting as well as properties of the used matured co-compost and irrigation waters were performed using classical methods as reported and explained by Cottenie *et al.*(1982); Page *et al.*(1982); Carter(1993); Rowell (1996); Tan(1993) and Burt(2004) as tabulated in all Tables in this work. Ionic strength (mmoles L⁻¹): was calculated using the following equation as explained by Tan(1993).

$$\text{Ionic strength (mmoles L}^{-1}\text{)} = \frac{1}{2} \sum_{i=1}^{i=n} M_i Z_i^2$$

Where: M_i = conc. of ion (i) in mmoles L⁻¹ and Z_i = charge of ion (i).

Additionally, undisturbed vertical cylindrical volumes of field-moist soil samples were gently obtained using cylindrical sharp edged core samplers for estimating soil physical properties and soil moisture constants using routine work analysis methods as reported and described by Garcia(1978); Klute(1986); Okalebo *et al.*(1993) and Reynolds 1993(a,b). Soil moisture constants (field capacity, permanent wilting point and available water capacity) were measured and calculated by means of pressure cooker and pressure membrane apparatus for measuring moisture contents at pressures of 0.33 and 15 bar according to Garcia(1978) and reported by Klute(1986). Bulk and particle (real) densities were estimated as described by Blake and While, field saturated hydraulic conductivity *in situ* was determined using constant head well permeameter method employing Guelph permeameter apparatus as mentioned by Reynolds(1993b). All soil obtained values were calculated on oven dry weight basis 105 C ° for 24 hours. Some chemical properties of the different irrigation water sources under consideration during the carrying out of pot and field experiment periods are listed in Table (3). The suitable experimental design was selected for both pot- and field experiments. The wheat pot experimental cross-sectional area was 0.0453 m², while maize experimental plot area was 4.5 m² (1.5x3 m). The experimental design was split-split plot arrangement with three replications. The main plots were devoted to three irrigation treatments as follows: Wet – treatments (light irrigation) after 30 % AWSMD from soil available water capacity; Medium – treatments (moderate irrigation) after 50 % AWSMD from soil available water capacity (50% AWSMD) and Dry treatments (heavy

irrigation) after 70% AWSMD from soil available water capacity (70% AWSMD). Wheat and maize plants were exposed to deficit irrigation and started directly after life watering irrigation (El-Mohayaa irrigation) for achieving the selected available soil moisture depletion levels under consideration. Detailed experimental obtained data about irrigation scheduling and the actual seasonal applied water for wheat and maize crops production cultivated in loamy sand soils subjected to soil moisture depletion regimes over the growing winter and summer season periods 2010/2011 and 2011 are tabulated in Tables (6 and 7). The sub-plots were assigned to five types of soil conditioners and their mixtures 1:1:1 (w/w). The conditioner treatments (w/w) were applied as follows: Control (without additions); Bentonite at application rates of 0.2 % and 0.3%. ; Co-compost at application rates of 0.3 % and 0.5 %; Mixture of Natural Raw Minerals (MNRM) at application rates of 0.2 % and 0.3 %; and the mixtures of the three previous conditioners in 1:1:1 ratio at rates of 0.233% and 0.367%. The soil conditioner treatments were randomly distributed in the three main plots.

Table (1): Initiative physico-chemical characteristics of the selected arable experimental sites under consideration located at Abou-Omera Al-Sharkeya village, Baltim district before planting

- * Site(1): Properties of disturbed and undisturbed surface soil (0-30 cm) just before collection and transportation for wheat crop cultivation in pot-experiment.
- * Site(2): Properties of disturbed and undisturbed surface soil just before cultivation of maize crop field –experiment.

These conditioner types are mixed well with soil during its preparation for cultivating wheat before sowing and incorporated into soil surface before plowing during soil service process and its preparation before maize planting. Sub sub plots were occupied with two application rates as follows: R_1 and R_2 were (low) minimum and (high) maximum recommended application rates respectively Mixture Natural Raw Minerals (MNRM) and bentonite were purchased from Al-Ahram company for mining, natural minerals (ores) and fertilizers. These materials are the new products from Al-Ahram Company for improving soil properties and fertility. The chemical analysis of these materials listed in Table (4). The analytical data of elemental oxides were kindly obtained from Al-Ahram company. These natural minerals were used as soil conditioners for wheat pot – and maize-field experiments.

Seeds of wheat plants (*Triticum aestivum*, Sakha 93 variety) were obtained from Crop Agronomy Research Department, Sakha Agriculture Research Station, Ministry of Agriculture and Land Reclamation. Wheat pot-experiment was conducted on experimental research area of Sakha Agriculture Research Station, Kafr El-Sheikh city. Pot experiment was performed using cylindrical perforated plastic pots having (mean internal diameter 24 cm and height 21 cm) under wire proof greenhouse conditions. Pot cross-sectional area was 0.0453 m^2 and its interval volume 9.504 liters. Composite loamy sand soil was collected and brought from fruit field (Site 1) located at Abou-Omera east village, Baltim district as mentioned before. Each pot contained 10 Kg soil on oven dry weight basis, wheat cultivation elongated 135 days. Throughout the wheat growth period, a freely drained water was collected from each plastic pot and reused again with irrigation water and also whenever it was necessary.

Grains of maize plants (*Zea mays, L*) three cross 321 variety were obtained from Maize Research Center, Agriculture Research Center, Ministry of Agriculture and Land Reclamation. Maize field- experiment was carried out on cultivated area of Abou-Omera east village (Site 2), Baltim district during the growing summer season period 2011 elongated 93 days. Total rented area = 3.5 kyrat = 612.5 m^2 and net cultivated area 405 m^2 (90 plots). Experimental plot area was 4.5 m^2 (1.5 x 3) and its weight 1046.25 kg loamy sand soil on oven dry weight basis.

Table (2): Soil moisture constants and its nutritional status of the selected experimental sites under consideration located at Abou-Omera Al-Sharkeya village, Baltim district before planting.

Soil variables	Obtained values	
	Site (1)* Pot-experiment	Site(2)** Field-experiment
Soil moisture constants		
Soil field capacity(S.F.C) %	17.0	18.0
Soil permanent wilting point(P.W.P) %	8.50	9.00
Soil available water capacity(A.W.C) %	8.50	9.00
Soil nutritional status		
Total organic-C %	0.174	0.232
Organic matter(O.M) %	0.298	0.400
Available macro-nutrients		
Available – N(K-sulphate extractable) $mgkg^{-1}$ soil	18.28	21.5
Available – P(NaHCO ₃ extractable) $mgkg^{-1}$ soil	7.62	8.90
Available – K(NH ₄ -acetate extractable) $mgkg^{-1}$ soil	50.15	53.5
Available micronutrients		
Available – Fe(DTPA extractable) $mgkg^{-1}$ soil	6.00	6.50
Available - Mn(DTPA extractable) $mgkg^{-1}$ soil	4.45	5.00
Available-Zn(DTPA extractable) $mgkg^{-1}$ soil	1.20	1.10
Available – Cu (DTPA extractable) $mgkg^{-1}$ soil	0.34	0.66

Notes : See feet notes of Table(1).

Aerobic / Thermophilic co-composting process was carried out at the experimental farm of Soil Improvement and Conservation Research Department, Sakha Agriculture Research Station during the summer growing season elongated five months from May 2010 to October 2010. Pyramidal piles(heaps) 2.5 × 2.5 × 1.5 m were built up under aerobic conditions. Different solid bio-wastes were used as substrates and augmented organically with farmyard manure (10 % w/w) as microbial organic activator as well as with urea , super phosphate and potassium sulfate as microbial chemical activators. The other certain additional materials were incorporated into for speeding up the conversion and improving the final product quality and as growth promoting substances, pH buffering agents and as bulking agents . The obtained chemical and physical characteristics of the used matured co-compost after co- composting process are listed in Table (5). This matured co-compost was used as soil conditioner.

Table (4): Chemical analysis of the used natural raw minerals and soil conditioners

Characteristics	Values	
	Bentonite	MNRM
Elemental oxides: %		
SiO ₂	55.9	39.36
TiO ₂	0.20	0.81
Al ₂ O ₃	20.0	7.68
Fe ₂ O ₃	0.70	4.05
MnO	0.001	0.67
MgO	0.65	3.20
CaO	2.70	15.07
Na ₂ O	1.76	1.95
K ₂ O	2.40	3.94
P ₂ O ₅	0.80	7.33
SO ₃	-	5.83
Loss on ignition	10.0	9.14
ECe dS m ⁻¹ (1:10 Bentonite-water extract(w/v))	1.82	
pH (1:2.5 bentonite-water suspension (w/v))	7.12	
Total soluble cations (meq L ⁻¹) (1:5 extracts)		
Ca ⁺²	0.79	
Mg ⁺²	0.27	
Na ⁺	1.95	
K ⁺	0.02	
Total soluble anions (meq L ⁻¹) (1:5 extracts)		
CO ₃ ⁼	-	
HCO ₃ ⁻	0.24	
Cl ⁻	1.59	
SO ₄ ⁼	1.06	
Cation exchange capacity, cmoles kg ⁻¹	59.13	
Calcium carbonate %	14.27	
Particle size distribution %		
Clay fraction	85.75	
Silt fraction	10.54	
Sand fraction	3.71	

Notes:

1- MNRM: Mixture of Natural Raw Minerals

2- The analytical results of the elemental oxides were kindly obtained from Al-Ahram company for mining and natural fertilizers.

Table (5): Chemical properties of the used co-compost directly after composting process

Characteristics	Values
Dry weight (kg m ⁻³)	650.0
Moisture content (%)	25.5
Odour and colour	Acceptable and dark
pH (1:10 compost-water suspension w/v)	7.16
EC (1:10 compost – water extraction w/v)	5.23
Total soluble salts(soil paste –water extraction 1:10)%	0.335
Saturation percentage % (g/100g	175.0
Total soluble salts (compost material)% (g/100g compost)	0.586
CEC (cmole kg ⁻¹)	64.34
Total organic – c %	25.5
Total organic matter %	43.96
C/N ratio	21.98
Total macro-nutrients %	
Total – nitrogen %	1.16
Total – phosphorus %	0.53
Total – potassium %	0.37
Available macro-nutrients (mg kg compost)	
Available – N (potassium sulfate)	100
Available – P (0.5 M NaHCO ₃ - pH 8.5)	50
Available – K (ammonium acetate pH 7)	85
Available micro-nutrients (mg kg compost)	
Available – Fe	450
Available – Mn	100
Available – Zn	35
Available – Cu	135
Total micro-nutrients (mg kg compost)	
Total –Fe	753
Total – Mn	361
Total – Zn	297
Total – Cu	168
Available heavy metals (mg kg compost)	
Available – cd	13.2
Available – Ni	62.7
Available – pb	120

Irrigation water supply:

Irrigation water supply and number of irrigations were limited according to the levels of soil moisture depletion regimes. Consequently, soil moisture content at demand depletion levels determines the timing of irrigation. Soil moisture content directly before irrigation at which calculated applied water must be added immediately for arriving at soil field capacity was measured *in situ* using TDR apparatus (Time Domain Reflectometer). Magnitude of irrigation applied water were calculated using the following soil moisture depletion equation as reported by (Israelson and Hansen, 1962) during wheat and maize growing season periods.

$$Q = \frac{SFC - CMC}{100} \times Bd \times D \times A$$

Where: Q = Quantity of applied water m³ pot⁻¹ /irrigate for pot-experiment, and m³ plot⁻¹ /irrigate for field –experiment ;SFC = Soil field capacity (%) in percent by volume; CMC = Soil moisture content just before irrigation using TDR apparatus; Bd = Soil bulk density Mg m⁻³ ; D = Soil depth (m), effective root depth or soil depth required to be irrigated; and A = pot or plot experimental area (m²) that would be irrigated. With respect to maize field water measurements, the magnitude of planting and life watering irrigates were measured and applied using cutthroat flume(20 ×90 cm)according to Early(1975).

A common NPK-fertilization was applied to the soil active root zone during the wheat and maize growing seasons according to the recommended doses of Ministry of Agriculture for wheat and maize crops under sandy soil conditions.

At harvesting time after wheat and maize plants full maturity, biomass grains and straw yields were fairly hand pulled and collected from each wheat pot experiment as well as from inner two rows of central area of maize plots. Some agronomical characteristics of these cereal crops and their productions such as (biomass grains, straw, biological yields and weight of 1000 wheat grains and 100 maize grains) as well as yield vegetative features as affected by soil conditioner types, their mixtures and application rates under irrigation deficits in the studied sandy soil were weight, measured, estimated, recorded and calculated some other parameters. Harvest index(%) was calculated as follows:

HI % = biomass grains yield / biological crop yield × 100

Statistical analysis:

Analysis of variance was done according to (Snedecor and Cochran,1976) using the Irristat software, version 4.1 according to Biometrics Unit,1998, IRRI(1998).

RESULTS AND DISCUSSION

Effect of applying soil conditioners under irrigation deficits on soil properties after cereal crops harvesting

Chemical characteristics of soil suspensions and extractions:

Concerning the effect of applying natural soil conditioner types, their mixtures(1:1:1) and application rates in sandy soils subjected to soil moisture depletion regimes (30%, 50 % and 70 %) from their available water capacities after wheat and maize crops harvesting on soil salinity (pH, EC dS m⁻¹, and ionic strength mmole L⁻¹) and hazard sodium parameters (SAR and SSP %) are listed in Tables (8.1 and 8.2). Generally, the analytical chemicals results listed in aforementioned tables illustrate that values of these chemical parameters in the studied soil on the average of other studied parameters (irrigation treatments and conditioner application rates) were markedly increased due to the application of soil conditioner types in comparison with control values (without additions). These increase could be arranged in the following descending order as follow: Mixtures (1:1:1) > MNRM > Bentonite > Compost >> control .

It is obviously that, the highest values were achieved as a result of mixtures application, meanwhile, the lowest values were obtained by dressing the compost treatment. However, the highest pH values were recorded at MNRM application under wheat pot experiment. Meanwhile, the lowest values of I.S were obtained at the application of bentonite under field-maize experiment.

On the other hand, these studied chemical properties, on the average of the other studied parameters (conditioner types, and their application rates), were markedly increased with increasing the depletion regimes from its soil available water capacity. Where, the highest values of these chemical parameters were achieved under dry treatment (70 % AWSMD), meanwhile, the lowest values were recorded under wet treatment (30 % AWSMD). Medium treatment had the moderate values between wet and dry treatments. The analytical obtained increments could be rearranged in the following ascending order: Wet –treatment (30 % AWSMD) < Medium – treatment (50 % AWSMD) < Dry – treatment (70 % AWSMD). This could be attributed to the dilution effect , since, salt concentration was decreased with increasing irrigation applied water. As delineated in Tables (8.1 and 8.2), obtained values of the studied chemical properties on the average of the other studied parameters (condition treatments and irrigation regime treatments) were higher under high application rate (R_2) rather than under low application rate (R_1) at the same conditions. Data listed in Table (8.2) reveal also that under maize field experiment , the studied chemical properties were increased with adding soil conditioner types and their mixtures (1:1:1) on the overall average of the other studied parameters (irrigation treatments and conditioners application rates) in comparison with their control –values (without additions) at the same conditions. These parameters mannered the following descending order: Mixtures (1:1:1) > MNRM > Bentonite > Co-compost > Control. Generally, the obtained values of chemical properties after wheat crop harvesting were higher than those obtained after maize crop harvesting.

Soil physical properties:

Concerning the effect of natural soil conditioner types, their mixtures 1:1:1 (w/w) and application rates under soil moisture depletion levels from its available water capacity on soil physical properties after wheat and maize crops harvesting are presented in Tables(9.1 and 9.2). Generally, the results collected in Table(9.1) show that, on average of other studied parameters (irrigation treatments and conditioner application rates), that saturated hydraulic conductivity SHC ($m\ day^{-1}$) and bulk density D_b ($Mg\ m^{-3}$) were markedly decreased as a result of adding soil conditioner types and their mixtures (1:1:1) in the studied soil after wheat crop harvesting in comparison with their control values. The magnitude of these decrements which less the control values were depended upon the types of these conditioners. It is clearly that, the lowest values were achieved and accompanied with applying conditioner mixtures 1:1:1 (w/w) treatment. Meanwhile, the highest values were obtained with the dressing compost conditioner in comparison with their control values. These decrements could be arranged in the following descending order as follows: Mixtures 1:1:1 > MNRM > Bentonite > compost > control. Oppositely, as demonstrated in the above mentioned Table,

soil porosity values were mannered the opposite trend, where such values were increased over the control –values with adding soil conditioners and had the following sequence:

Mixtures (1:1:1) < MNRM < Bentonite < Compost < Control

Commonly, the obtained values of all studied physical properties on the average of the other studied parameters (conditioner treatments and irrigation treatments) under the high application rate (R_2) were lower than those obtained under the low application rate (R_1). On the other hand, it was clearly apparent that, SHC values were gradually increased with increasing water irrigation deficits. However, D_b and ρ_r had the opposite trend, which decreased with increasing soil moisture depletion levels for its available water capacity. SHC – increments as well as D_b and ρ_r decrements with increasing moisture depletion levels could be arranged in the following descending order as follows:

Wet – treatment (30% AWSMD) > Medium – treatment (50% AWSMD) > Dry – treatment (70% AWSMD)

In respect of maize field experiment, data demonstrated in Table (9.2) show on average of irrigation treatments that, application of all natural soil conditioner types and their mixtures (1:1:1) resulted in decreasing saturated hydraulic conductivity and soil bulk density, as well as led to increasing total porosity in comparison with their control values at the same conditions. It was clear that, conditioner mixtures (1:1:1) realized the lowest values of SHC and ρ_r besides the highest values of D_b in comparison with their control values. However, application of co-compost treatment mannered the opposite trend, which gave the highest values of SHC and ρ_r in addition to the lowest values of D_b . Generally, it was apparent from the results that, on average of irrigation treatments and conditioner treatments that, adding high application rates lowered gradually SHC and D_b and raised the values of ρ_r . In this direction, the low application rates led to the opposite trend. Commonly, analytical data listed in Tables (9.1 and 9.2) illustrate, on average of all other studied parameters (conditioner treatments and application rates) that, SHC and ρ_r values were gradually increased with increasing soil moisture depletion levels from its available water capacity. However, D_b values were decreased with increasing irrigation deficits.

Soil moisture constants:

As concerns, field capacity (SFC %), permanent wilting (PWP %) and available water capacity (AWC%) values of the studied sandy soils which reflect their soil water holding capacity after wheat and maize crops harvesting as affected by soil conditioner types, their mixtures and application rates under soil moisture depletion regimes, are presented in Tables (10.1 and 10.2) respectively. The obtained results, demonstrate, on average of the other studied parameters (irrigation treatments and conditioner –application rates) that, these soil moisture constants were obviously increased by applying soil conditioners and their mixtures (1:1:1) in comparison with their control values at the same conditions.

Generally, these increments over the controls in studied soil after wheat crop harvesting could be arranged in the following sequence as follows:

Mixtures (1:1:1) > MNRM > Bentonite > Compost > Control

Regarding maize field experiment, data listed in Table (10.2) reveal, on average of other studied parameters that, values of soil moisture constants also were increased in comparison with their control values at the same conditions by applying soil conditioner types and their mixtures (1:1:1). These increments of all soil moisture constants in studied soil after maize crop harvesting could be rearranged in the following sequence as follows :

Compost > Mixtures (1:1:1) > MNRM > Bentonite > Control

High conditioners application rate generally was realized higher values than those obtained by low conditioners application rate. As delineated in Table (10.1) it was clearly apparent on average of the other studied parameters, that these water properties were gradually decreased with increasing soil moisture depletion levels for its available water capacity. The highest values were achieved under wet – treatment (30% AWSMD) while, the lowest values were given by dry – treatment (70% AWSMD). Medium – treatment (50% AWSMD) recorded the moderate values of these soil moisture constants between wet(30 % AWSMD) and dry(70 % AWSMD)treatments.

Analytical results show that applying conditioner-mixtures (1:1:1) show its superiority over all other conditioner types under wheat –pot experiments. Meanwhile, applying compost treatment show its superiority over all other conditioner types under maize field experiment.

Effect of applying soil conditioners under irrigation deficits on soil macronutrients content:

Concerning residual contents (concentrations) of (N and P) after wheat and maize crops harvesting as affected by soil conditioner types, their mixtures and application rates subjected to irrigation deficits were listed in Table (11.1). Analytical results listed in this table reveal that, on average of other studied parameters (irrigation treatment and conditioner application rates), that N and P macronutrients content in studied sandy soils were increased by adding soil conditioner types and their mixtures (1:1:1) after wheat and maize crops harvesting in comparison with their control values.

The highest values of residual N and P were achieved and accompanied by mixtures (1:1:1) treatment. Meanwhile, dressing compost treatment gave lower values than those obtained by applying mixtures treatment in comparison with their control after pot wheat experiment at the same conditions.

Oppositely, residual N and P macro nutrients in the studied soil after maize crop harvesting behaved the opposite trend, where, the highest contents of these nutrients were achieved by applying compost treatment. However, the dressing conditioner mixtures gave values lesser than those obtained by adding compost treatment. The lowest values were absolutely obtained in the studied soil by adding bentonite treatment after either wheat or maize crop harvesting. These increments of the residual N and P

macronutrients content after wheat crop harvesting could be arranged in the following sequence as follows:

Mixtures treatment > MNRM > Compost > Bentonite > Control. Moreover, on average of other studied parameters, obtained residual values of these macronutrients under high rate realized slightly values higher than those obtained under low application rate for either wheat or maize crops harvesting.

It was clearly apparent as delineated in Table (11.1) on average of other parameters, that content of macronutrients (N and P) in the studied sandy soil after wheat and maize crops harvesting under medium –treatment (50% AWSMD) were realized the highest values then under wet –treatment, while, the lowest values were obtained under dry-treatment. Residual contents of N and P macronutrients after maize crops harvesting could be also arranged in the following descending order: Compost treatment > MNRM > Mixtures > Bentonite > Control. Residual N and P macronutrients in studied sandy soils after wheat and maize crop harvesting could be arranged in the following descending order as: Medium –treatment (50%AWSMD) > Wet –treatment (30% AWSMD) > Dry-treatment (70% AWSMD). From the abovementioned results, it could be concluded that, on average other studied parameters, residual N and P macronutrients content in studied soils after crops harvesting means low exhausting nutrients from soils during the growing seasons of wheat and maize crops.

Effect of dressing soil conditioners under moisture depletion regimes on agronomical crops production

Biomass grains and straw yields:

Regarding wheat crop pot experiment after full maturity, data listed in Tables (12.1, 12.2 and 12.3) demonstrate, on average other studied parameters (irrigation treatments and conditioner application rates), that application of soil conditioner types and their mixtures (1:1:1) resulted in significantly increasing wheat biomass grains yield, thousand grains weight, harvesting index and biomass straw yield in comparison with their control values at the same conditions.

Moreover, on average of the other studied parameters, the values of these agronomical features under high application rate (R_2) were higher than those obtained under low application rate (R_1). The increments of these agronomical traits could be arranged in the following descending order as: Mixtures (1:1:1) > MNRM > Bentonite > Compost > Control. So, the highest values of these agronomical features in studied sandy soils were achieved and accompanied by applying conditioner – mixtures treatment. Meanwhile, applying compost treatment gave the lowest values. However, the analytical obtained data, on average of the other studied parameters (conditioner treatments and their application rates) reveal that all agronomical features with the exception of harvest index were gradually decreased with increasing soil moisture depletion levels from its available water capacity. The magnitude of these decrements could be arranged in the following descending order as: Wet-treatment (30% AWSMD) > Medium –treatment (50% AWSMD) > Dry –treatment (70 % AWSMD).

However, harvest index (%) was slightly increased with increasing irrigation deficits. Therefore, these increments could be arranged in the following sequence as: Dry-treatment (70 % AWSMD) > Medium-treatment (50% AWSMD)>Wet-treatment (30% AWSMD).

Concerning field maize crop after full maturity, tabulated data show on average other parameters (irrigation treatments and conditioner application rates) that, dressing soil conditioner types and their mixtures 1:1:1 (w/w) led to increasing significantly biomass maize grains yield, hundred grains weight, harvesting index and biomass straw yield in comparison with their control values at the same conditions with the exception of harvest index. Moreover, the values of these agronomical features under high application rate (R_2) gave higher values than those obtained under low application rate (R_1) with exception of H.I which behaved the opposite trend i.e. $R_1 > R_2$. On the other hand, the obtained data under wet treatment (30% AWSMD) on average of the other studied parameters (conditioners treatment) that compost treatment gave the highest values of these studied parameters except harvest index which had an opposite trend, where the highest values were obtained by adding their mixtures in comparison with compost application. It could be due to increasing straw yield of compost treatment as comparison with under their mixtures treatment. The lowest obtained values were generally recorded under bentonite treatment. However, such parameters under medium and dry treatments behaved the opposite trend in comparison with wet treatment, since the maximum values of biomass grains yield, 100 grain weight and harvest index were realized under mixtures treatment. Meanwhile, the biomass straw yield under compost treatment was higher than those obtained under mixture treatment. Generally, mean values of these parameters under medium treatment were higher than those obtained under wet and dry treatments and behaved the following order: Medium > Wet > Dry. Such results were obtained by Abdel-Reheem and Hassan (2011) , they found that the highest values of wheat water productivity and yield were achieved when irrigation at 50 % depletion from available water , compared to 70 % and 40 % depletion in the loamy soils. confirming this conclusion, similar responses of maize crop production under field conditions was also reported by Khalifa (2013), who stated that, irrigation at 50 % SMD gave the highest values of yield and its components of maize crop.

Crop yield components:

As regards to wheat and maize crops vegetative features after full maturity as affected by soil conditioner types, their mixtures and application rates subjected to different levels of irrigation deficits were presented in Tables (12.4 and 12.5). Obtained data shown in Table (12.4) reveal on average of the other studied parameters that, the following wheat crop vegetative features (plant height, number of tillers/ spike; number of spikelets/spike; spike length and panicle mean weight were markedly increased as a result of adding soil conditioners and their mixtures 1:1:1 (w/w). These increments could be arranged in the following descending order:

Mixtures 1:1:1 treatment > MNRM treatment > Bentonite treatment > Compost treatment > Control. Furthermore, the mean values of these crop vegetative features under high application rate (R_2) were higher than those obtained under low application rate (R_1) i.e. $R_2 > R_1$. On the other hand, on average of the other parameters, all these features were gradually decreased with increasing soil moisture depletion levels from its available water capacity. Consequently, these obtained decrements could be arranged in the following order as: Dry –treatment (70% AWSMD) < Medium –treatment (50% AWSMD) < Wet-treatment (30% AWSMD).

With respect to maize field crop vegetative features after full maturity, data were listed in Table (12.5). Obtained data show, on average of the other parameters, that some maize crop vegetative features such as plant height, ear weight, and ear length were obviously increased by adding soil conditioners and their mixtures 1:1:1 (w/w). High application rate (R_2) recorded values higher than those obtained by low application rate (R_1). So, under field experiment, compost treatment gave the high values in comparison with mixtures treatment at the same conditions, bentonite treatment realized the lowest values. Respecting irrigation regimes under field experiment, data show on average of the other studied parameters that medium –treatment (50% AWSMD) gave the highest values followed by wet –treatment (30% AWSMD), while, dry –treatment recorded the lowest values.

REFERENCES

- Abdel-Rheem , H.A. and A.F. Hassan(2011). Reducing of water use by water stress regime on some main field crops (wheat , soybean and corn). J. Soil. Sci . and Agric.Eng.,Mansoura Univ.,Vol.2(6):635-648.
- Blake, G.R. and K.H. Hartge (1986 a). Bulk density. Pages 363 – 375. In A. Klute (ed.): Methods of Soil Analysis. Part 1 (2nd Ed.): Physical and Mineralogical Methods. ASA, Inc., SSSA, Inc., Publisher Madison, Wisconsin, USA.
- Blake, G.R and K.H. Hartage (1986 b). Particle density. Pages 377 – 382. In A. Klute (ed.): Methods of Soil Analysis. Part 1 (2nd Ed.): Physical and Mineralogical Methods. ASA, Inc., SSSA, Inc., Publisher Madison, Wisconsin, USA.
- Burt,R.(2004). "Soil Survey Laboratory Methods Manual". USDA – NRCS, Linoln, Nebraska.
- Carter, M.R.(ed.)(1993). Soil Sampling and Methods of Analysis. Canadian Society of Soil Science. Lewis Publishers., Boca Raton, London, Tokyo.
- Carter, M.R. and B.C.Ball(1993). Soil porosity. Pages 581 – 588. In M.R. Carter (ed.): Soil Sampling and Methods of Analysis. Canadian Society of Soil Science. Lewis Publishers, Boca Raton, London, Tokyo.
- Cottenie, A.; P.M. Verloo; L.Kiekens ; G.Velghe and R. Camerlynck(1982). "Chemical Analysis of Plant and Soils". Lab.Anal. and Agrochem. State Univ., Gent, Belgium.

- Early, A.C. (1975). Irrigation scheduling for wheat in Punjab. Centosci Prog. Optimum use of water in agriculture, Rpt, 17, Layllpur, Pakistan, 3-5, March, Pages 115 – 127.
- El-Hady, O.A. and A.F. El-Sherif (1988). Egyptian bentonite deposits as soil conditioners. II: Hydro physical characteristics and mechanical strength of sandy soils treated with bentonite. Egypt. J. Soil Sci., 28: 215-223.
- Garcia, I. (1978). Soil Water Engineering Laboratory Manual. Department of Agricultural and Chemical Engineering. Colorado State University, Fort Collins, Colorado, USA.
- IRRI (1998). Using Irristat Software in Statistical Analysis. Manual, Biometrics Unit. Filipinas.
- Israelson, O.W. and V.E. Hansen (1962). Irrigation Principles and Practices (3rd Ed.). John Willey & Sons Inc., New York.
- Khalifa, R.M. (2013). Water requirements of maize and sugar beet crops affected by soil moisture depletion and water table level. M.Sc. Thesis, Soil Sci. Dept. Fac. of Agric., Kafrelsheikh Univ.
- Klute, A. (ed.) (1986). Methods of Soil Analysis, Part 1 (2nd Ed.): Physical and Mineralogical Methods. ASA, Inc., SSSA, Inc., Publisher, Madison, Wisconsin, USA.
- Okalebo, J.R.; K. W. Gathua and P.L. Woomer (1993). Laboratory Methods of Soil and Plants Analysis: A Working Manual. TSBF Program, Soil Sci. Soc. of East Africa. Technical Publication. No.1, Unesco, Rosta, Kenya.
- Omran, W.M. (2005). Maize yield response to available soil moisture. Minufiya J. Agric., Res., Egypt, 30(4): 1257 – 1268.
- Page, A.L.; R.H. Miller and D.R. Keeney (eds.) (1982). Methods of Soil Analysis. Part 2 (2nd Ed.): Chemical and Microbiological Properties. ASA Inc., SSSA, Inc. Publisher, Madison, Wisconsin, USA.
- Reynolds, W.D. (1993 a). Saturated hydraulic conductivity: Laboratory measurement. Pages 589 – 598. In M. R. Carter (ed.): Soil Sampling and Methods of Analysis. Canadian Society of Soil Science. Lewis Publishers, Boca Raton, London, Tokyo.
- Reynolds, W.D. (1993 b). Saturated hydraulic conductivity: Field measurement. Pages, 599 – 613. In M. R. Carter (ed.): Soil Sampling and Methods of Analysis. Canadian Society of Soil Science. Lewis Publishers, Boca Raton, London, Tokyo.
- Rowell, D.L. (1996). Soil Science: Methods and Applications. Longman, UK.
- Seif, S.A., S.A.H. Allam; M.E. El-Emery and A.E.M. El-Galfy (2005). Effect of soil moisture depletion on growth, yield and yield components of some maize varieties. Annals of Agric. Sci., Moshtohor, 43(1): 25 – 58.
- Snedecor, G.W. and W.G. Cochran (1976). Statistical Methods 6th (ed.), Iowa State Univ. press, Iowa, USA.
- Tan, K.H. (1993). Principles of Soil Chemistry. (2nd Ed.). Page 62. Marcel Dekker Inc., New York.
- van Wambeke, A. (1992). Soils of the Tropics – Properties and Appraisal. McGraw Hill, Inc., New York, USA.

Várallyay G.Y. (2005): Role of soil multifunctionality in future sustainable agricultural development. Acta Agronomica Academiae Scientiarum Hungaricae, 51:109–124.

فاعلية إضافة محسنات التربة الطبيعية تحت مستويات استنزاف رطوبي على خواص التربة الرملية وإنتاجية محاصيل الحبوب

محمد على محمد القماح¹، مروة جمال محمد على²، محمود احمد ابوالسعود² وسمير على مشالي¹

1- قسم الأراضي والمياه ، كلية الزراعة ، جامعة كفر الشيخ ، كفر الشيخ، مصر.
2- معهد بحوث الأراضي والمياه والبيئة، محطة البحوث الزراعية بسخا، كفر الشيخ، مصر

الهدف من البحث هو دراسة تأثير إضافة محسنات التربة الطبيعية ومخاليطها تحت مستويات من الاجهاد الرطوبي على خواص التربة الرملية وإنتاجية بعض محاصيل الحبوب.

لهذا الغرض اقيمت تجربتان في موقعان يمثلان الأراضي الرملية بمصر بقرية ابوعميرة الشرقية، مركز بلطيم، محافظة كفر الشيخ. الموقع الاول حقل فاكهة اخذت منه عينات سطحية مركبة مثارة واخرى غير مثارة، تم دراسة صفاتها الكيميائية والفيزيائية والمائية لإجراء تجربة قصارى لزراعة القمح بمحطة البحوث الزراعية بسخا خلال الموسم الشتوي 2011/2010 استمرت 135 يوم، والموقع الثاني قريب من الاول تم استخدامه لإجراء تجربة حقلية لزراعة الذرة في الموسم الصيفي 2011 استمرت 93 يوم. اضيفت الاسمدة الازوتية والفوسفاتية والبوتاسية الموصى بها، واتبع تصميم القطع منشقة المنشقة في ثلاث مكررات (90 وحدة تجريبية).

تضمنت المتغيرات الدراسية: اربعة معاملات للمحسنات الطبيعية (مخلوط خام المعادن الطبيعية، الكميوست، البنتونيت و مخلوط شامل من هذه المحسنات بنسبة 1:1:1)، معدلين إضافة أعلى وأقل من الموصى به، ثلاث مستويات من الاستنزاف الرطوبي من الماء الميسر في التربة (30 % ، 50 % ، و70%).

اهم النتائج الدراسية المتحصل عليها يمكن تلخيصها فيما يلي:-

1- ازدياد قيم صفات الملوحة ورقم تفاعل التربة، التوصيل الكهربى والقوة الايونية لمعلقات ومستخلصات التربة (1:5) وكذلك قيم محددات ضرر الصوديوم (SAR, SSP%) بعد حصاد القمح والذرة نتيجة لإضافة المحسنات الطبيعية للتربة الرملية كمتوسط لتأثير باقي المتغيرات الدراسية مقارنة بالكنترول. وقد حققت إضافة المخلوط الشامل (1:1:1) أعلى القيم مقارنة بالكميوست بالتربة بعد حصاد القمح والذرة. وقد سجل معدل الإضافة الثاني قيم أعلى من معدل الإضافة الأول. وازدادت قيم هذه المتغيرات الكيميائية كقيمة متوسطة لتأثير باقي المتغيرات بزيادة مستوى الاستنزاف الرطوبي وقد اعطت المعادلة المعتدلة AWSMD % 50 قيم وسطية بين المعاملة الرطبة AWSMD % 30، والمعاملة الجافة AWSMD % 70 .

2- تحسن في الخواص الطبيعية للتربة بانخفاض قيم الكثافة الظاهرية والتوصيل الهيدروليكي وارتفاع قيم المسامية الكلية للتربة مقارنة بالكنترول وتأخذ قيم الانخفاض وزيادة الترتيب التصاعدي التالي:

Mixtures (1:1:1) > MNRM > Compost > Bentonite > Control

وقد حقق معدل الإضافة الأعلى انخفاضاً في قيم H_C , D_b وارتفاعاً في قيم المسامية الكلية مقارنة بمعدل الإضافة الأدنى، وعموماً في تجارب الأصص للقمح حققت إضافة المخلوط الشامل أفضل تحسن في الخواص الطبيعية في حين أن تجارب الذرة الحقلية حققت إضافة الكميوست أفضل النتائج.

وقد ازدادت قيم الخواص الطبيعية بزيادة مستوى الاجهاد المائي فقد حققت المعاملة المعتدلة AWSMD % 50 قيم وسطية بين المعاملة الرطبة والجافة.

3- تحسنت قدرة التربة على الاحتفاظ بالماء متمثلة في زيادة ثوابت رطوبتها الارضية (السعة الحقلية ، نقطة الذبول والماء الميسر) بإضافة المحسنات الطبيعية ، فقد حققت إضافة المخلوط الشامل قيم أعلى لهذه الخواص المائية تحت ظروف تجارب الأصص ، في حين حققت إضافة الكميوست أعلى القيم تحت ظروف التجارب الحقلية. واطهرت النتائج أيضاً زيادة قيم هذه الثوابت عند إضافة المعدل الأعلى مقارنة بالمعدل الأدنى. ومن ناحية أخرى انخفضت قيم هذه الثوابت بزيادة مستوى الاستنزاف الرطوبي فلي تجربة الأصص واخذت الاتجاه العكسي في التجارب الحقلية . عموماً المعاملة المعتدلة AWSMD % 50 اعطت نتائج وسطية بين المبتلة والجافة.

4- زيادة محتوى التربة من العناصر الغذائية الكبرى (النيتروجين والفوسفور) المتبقية بعد حصاد محصولي القمح والذرة مقارنة بالكنترول نتيجة لإضافة المحسنات وقد حقق المخلوط الشامل أعلى تركيز متبقى لهذه العناصر في تجارب الأصص اما تحت ظروف التجارب الحقلية فكانت السيادة لإضافة الكميوست ، ومعدل الإضافة الأعلى حقق أعلى تركيز متبقى مقارنة بمعدل الإضافة الأدنى. معاملة الري المعتدلة ادت الى قيمة وسطية بين الرطبة والجافة.

5- زيادة مغنوية في محصول الحبوب والقش والبروتين ووزن وحدة الحبوب ، دليل الحصاد وكذلك مكونات محصولي القمح والذرة بعد تمام النضج نتيجة لإضافة المحسنات الطبيعية مقارنة بقمم الكنترول. وحقق معدل الإضافة الأعلى

افضل من معدل الاضافة الادنى، وكانت افضل معاملة هي المخلوط الشامل تحت ظروف تجربة الاصص ، في حين كانت معاملة الكمبوست لها السيادة تحت ظروف التجارب الحقلية. وعموما المعاملة المعتدلة اعطت نتائج مرضية وسيطة بين المبتلة والجافة

كلية الزراعة – جامعة المنصورة
مركز البحوث الزراعيه

أ.د / أحمد عبد القادر طه
أ.د / السيد عوض محمد

قام بتحكيم البحث

Table (6): Irrigation scheduling and actual seasonal applied water at different regular- intervals for wheat crop production cultivated in loamy sand soil under different soil moisture depletion regimes over the growing winter season period 2010/2011.

**Notes : 1- Total applied water/season = calculated applied water +planting and life watering irrigations.
2-Life watering irrigation = El-Mohayaa irrigation**

Table (7): Irrigation scheduling and actual seasonal applied water at different regular - intervals for maize crop production cultivated in loamy sand soil under different soil moisture depletion regimes over the growing summer season period 2012.

Irrigation scheduling	Wet – treatment 30 % AWSMD Light irrigation Short-intervals(every day)			Medium– treatment 50 % AWSMD Moderate irrigation Median -intervals(3 days)			Dry – treatment 70 % AWSMD Heavy irrigation Long -intervals(8 days)		
	Irrigation date	Applied water		Irrigation date	Applied water		Irrigation date	Applied water	
		Lplot ¹	m ³ fed ⁻¹		Lplot ¹	m ³ fed ⁻¹		Lplot ¹	m ³ fed ⁻¹
Planting irrigation (6.48 % SMC)	5/7/2012	241	224.6	5/7/2012	241	224.6	5/7/2012	241	224.6
Life watering irrigation (9.75 % SMC)	10/7/2012	172.6	161.12	10/7/2012	172.6	161.12	10/7/2012	172.6	161.12
Summation		413.6	385.7	Summation	413.6	385.7	Summation	413.6	385.7
Soil moisture content %	15.33 %SMC			13.5 % SMC			11.7 % SMC		
Regular- intervals	Short- intervals (every day)			Median- intervals(3 days)			Long- intervals (8 days)		
First irrigate	13/7/2012	56.5	52.73	15/7/2012	94.17	87.89	18/7/2012	244.82	228.5
Final irrigate	27/9/2012	56.5	52.73	25/9/2012	94.17	87.89	20/9/2012	244.82	228.5
Number of irrigates	77 irrigates plus planting and life watering irrigations			25 irrigates plus planting and life watering irrigations			9 irrigates plus planting and life watering irrigations		
Total irrigation period	86 days 5/7/2012 – 28/9/2012			86 days 5/7/2012-28/9/2012			86 days 5/7/2012-28/9/2012		
Maize harvesting time	October ,5,2012			October , 5, 2012			October , 5, 2012		
Maize growing season period	93 days			93 days			93 days		
Calculated applied water /season		4351	4060		2354	2197		2203	2057
Total applied water /season		4764	4446		2768	2583		2617	2442

Notes: Amounts of applied water for planting and life watering irrigations ($\text{ml plot}^{-1}/\text{irrigate}$) were measured using cutthroat flume (20x90) according to Early(1975).

A common NPK-fertilization was applied to the soil active root zone during -5

Table (8.1): Soil salinity, sodicity and the ionic strength of its extracts after wheat crop harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes.

Soil conditioner types and their mixtures	Conditioners application rates (w/w)	Soil moisture depletion levels from its available water capacity (AWSM D-levels)														
		Wet- treatment (30 % AWSMD) Light irrigation Short – intervals(3 days) 14.45 % SMC					Medium - treatment (50 % AWSMD) Moderate irrigation Median-intervals(6 days) 12.75 % SMC					Dry-treatment (70 % AWSMD) Heavy irrigation Long-intervals(9 days) 11.08 % SMC				
		Chemical parameters														
		pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹	pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹	pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹
SAR	SSP %			SAR	SSP %				SAR	SSP %						
Control	Without additions	7.80	0.32	2.86	64.71	3.64	7.60	0.35	2.86	63.36	4.45	7.40	0.44	3.26	63.13	5.36
Bentonite	low R1	8.06	0.58	3.76	62.50	8.63	8.12	0.62	3.85	62.69	8.78	8.30	0.75	4.14	62.82	9.88
	high R2	8.20	0.68	4.20	64.79	8.87	8.16	0.67	4.04	63.01	9.64	8.14	0.87	4.32	63.41	10.12
	Mean	8.13	0.63	3.98	63.64	8.60	8.14	0.645	3.94	62.85	9.21	8.22	0.81	4.23	63.11	10.00
Compost	low R1	7.83	0.54	3.24	60.00	6.76	8.12	0.68	4.04	63.01	7.63	8.10	0.73	3.51	61.20	8.24
	high R2	8.10	0.69	4.61	63.51	8.52	8.15	0.72	3.71	58.97	9.05	8.30	0.87	4.59	63.44	9.52
	Mean	7.96	0.61	3.92	61.76	7.64	8.13	0.70	3.87	60.99	8.34	8.20	0.80	4.05	62.32	8.88
MNRM	low R1	8.13	0.81	4.11	61.44	11.25	8.30	0.75	4.25	62.96	12.52	8.25	0.92	4.76	63.64	12.95
	high R2	8.21	0.97	4.63	62.38	12.67	8.18	0.74	4.24	63.29	13.28	8.40	0.94	4.77	63.37	14.72
	Mean	8.17	0.89	4.37	61.91	11.96	8.24	0.745	4.24	63.13	12.90	8.32	0.93	4.765	63.50	13.84
Their mixtures (1:1:1)	low R1	8.15	0.86	4.53	63.04	11.53	8.15	1.20	5.47	64.06	13.65	8.20	1.01	5.00	63.89	14.66
	high R2	8.15	1.13	4.29	63.64	13.36	8.20	1.12	5.17	63.33	14.95	8.22	1.07	5.09	63.48	15.28
	Mean	8.15	0.995	4.41	63.33	12.45	8.17	1.16	5.32	63.69	14.30	8.21	1.04	5.05	63.68	14.97
Average	low R1	8.04	0.698	3.90	61.74	9.54	8.17	0.813	4.40	63.18	10.65	8.21	0.86	4.58	63.47	11.42
	high R2	8.17	0.867	4.43	63.58	10.86	8.17	0.813	4.29	62.15	11.73	8.28	0.93	4.69	63.43	12.43
	Overall mean	8.10 3	0.781	4.17	62.66	10.18	8.17	0.813	4.345	62.67	11.19	8.24	0.895	4.64	63.45	11.93

Notes : 1 - Each value is a mean of

3 replications and all obtained values were calculated on oven dry weight basis at 105 C° for 24 hours.

2- Cylindrical plastic pots cross-sectional area (0.0453 m²) containing 10 kg loamy sand soil on oven dry weight irrigated with tap water

3-Winter growing season period of wheat crop elongated 135 days

4- Analytical data were determined and calculated using 1:5 soil water extracts (except pH)

5-Wet – treatment (light irrigation) : 30 % AWSMD for short –intervals (3 days)and actual seasonal applied water was 1107.4 m³ fed⁻¹(11.94 Lpot⁻¹)

6-Medium – treatment(moderate irrigation) : 50 % AWSMD for median –intervals (6 days)and actual seasonal applied water was 998.1 m³ fed⁻¹(10.76 Lpot⁻¹)

7-Dry – treatment(heavy irrigation):70 % AWSMD for long –intervals (9 days)and actual seasonal applied water was 972.4 m³ fed⁻¹(10.49L pot⁻¹).

8- SMC represents soil moisture content (%) directly before irrigation at which calculated water applied must be applied immediately for arriving its field capacity.

Soil conditioner types and their mixtures	Conditioners application rates (w/w)		Soil moisture depletion levels from its available water capacity (AWSM D-levels)														
			Wet- treatment (30 % AWSMD) Light irrigation Short – intervals(every day)				Medium - treatment (50 % AWSMD) Moderate irrigation Median-intervals (3 days)				Dry-treatment (70 % AWSMD) Heavy irrigation Long-intervals (8 days)						
			15.33 % SMC				13.5 % SMC				11.7 % SMC						
			Chemical parameters														
pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹	pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹	pH	EC dSm ⁻¹	H.S. parameters		I.S mmol L ⁻¹			
		SAR	SSP %				SAR	SSP %				SAR	SSP %				
Control	Without additions		7.84	0.35	2.69	64.52	3.58	7.64	0.38	2.58	62.50	4.46	7.42	0.46	2.17	54.05	5.35
Bentonite	low	R1	8.10	0.66	4.20	65.63	7.65	8.00	0.68	4.46	58.46	8.25	8.13	0.83	4.80	60.85	8.26
	high	R2	8.24	0.68	3.90	64.52	7.96	8.50	0.70	4.91	71.68	9.37	8.14	0.97	4.85	62.00	10.05
Mean			8.17	0.67	4.07	65.07	7.81	8.25	0.69	4.68	65.09	8.81	8.24	0.90	4.83	61.43	9.15
Compost	low	R1	7.80	0.57	3.25	59.50	7.65	8.17	0.53	3.44	53.06	8.00	7.80	0.82	3.41	55.80	9.02
	high	R2	8.19	0.69	4.20	62.63	8.36	8.19	0.61	4.56	58.80	8.58	8.50	0.90	4.72	60.81	9.68
Mean			7.99	0.63	3.73	61.06	8.01	8.18	0.57	4.00	55.94	8.29	8.18	0.86	4.06	58.30	9.35

MNRM	low	R1	8.10	0.85	4.15	61.05	10.13	8.23	0.74	3.38	57.97	10.08	8.05	0.92	4.09	59.78	10.62
	high	R2	8.16	0.97	4.32	62.50	12.45	8.28	0.75	4.66	67.56	11.36	8.50	0.94	4.66	64.44	11.40
Mean			8.13	0.91	4.24	61.78	11.29	8.25	0.745	4.02	62.77	10.72	8.30	0.93	4.38	62.11	11.01
Their mixtures (1:1:1)	low	R1	8.18	0.88	5.51	68.83	10.95	8.15	1.20	5.88	64.79	11.50	8.08	1.11	4.35	64.62	12.65
	high	R2	8.35	1.50	6.99	69.57	12.66	8.39	1.90	5.92	67.77	12.54	8.60	1.17	5.45	67.33	14.24
Mean			8.265	1.19	6.25	69.20	11.81	8.27	1.55	5.90	66.28	12.02	8.34	1.16	5.89	65.96	13.45
Average	low	R1	8.05	0.74	4.28	63.75	9.10	8.14	0.79	4.29	58.57	9.46	8.07	0.90	4.16	60.26	10.39
	high	R2	8.23	0.96	4.85	64.79	10.36	8.33	0.99	5.01	66.47	10.45	8.45	0.96	5.42	63.14	11.34
Overall mean			8.139	0.85	4.56	64.27	9.73	8.237	0.89	4.65	62.57	9.96	8.265	0.963	4.79	61.70	10.86

Table(8.2): Soil salinity, sodicity and the ionic strength of its extracts after maize crop harvesting as affected by natural soil conditioner types , their mixtures and application rates under different soil moisture depletion regimes

- Notes:**
- 1- Each value is a mean of 3 replications and all obtained values were calculated on oven dry weight basis at 105 C° for 24 hours.
 - 2- Analytical data were determined and calculated using 1:5 soil water extraction (except pH).
 - 3- Field plot sectional –area 4.5 m² and its weight 1046.25 kg loamy sand soil on oven dry irrigated with drainage water (Kotshner).
 - 4- Summer growing season period of maize crop elongated 93 days
 - 5- Wet – treatment (light irrigation): 30 % AWSMD for short –intervals (every day)and actual seasonal applied water was 4446 m³ fed⁻¹(4764 Lplot⁻¹)
 - 6- Medium – treatment (moderate irrigation): 50 % AWSMD for median –intervals (3days)and actual seasonal applied water was 2583 m³ fed⁻¹(2768 L plot⁻¹)
 - 7- Dry – treatment (heavy irrigation): 70 % AWSMD for long –intervals (8 days)and actual seasonal applied water was 2442 m³ fed⁻¹(2617L plot⁻¹).

8- SMC represents soil moisture content (%) directly before irrigation at which calculated applied water must be done immediately to arrive its field capacity.

Table (9.1): Saturated hydraulic conductivity, bulk density and total porosity of studied soils after wheat crop harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes .

Soil conditioner types and their mixtures	Conditioner application rates (w/w)		Soil moisture depletion levels from its available water capacity(AWSMD-levels)								
			Wet - treatment (30 % AWSMD) Light irrigation Short – intervals (3 days)			Medium - treatment (50 % AWSMD) Moderate irrigation Median – intervals (6 days)			Dry -treatment (70 % AWSMD) Heavy irrigation Long-intervals (9 days)		
			14.45 % SMC			12.75 % SMC			11.08 % SMC		
			Physical parameters								
		SHC m/day	D _b Mg /m ³	ρ _T %	SHC m/day	D _b Mg /m ³	ρ _T %	SHC m/day	D _b Mg /m ³	ρ _T %	
Control	Without additions		2.48	1.55	41.51	2.54	1.56	41.13	2.47	1.57	40.7
Bentonite	low	R1	2.43	1.43	46.00	2.52	1.45	45.28	2.55	1.42	46.42
	high	R2	2.41	1.45	45.28	2.40	1.41	46.79	2.43	1.40	47.17
Mean			2.42	1.44	45.66	2.46	1.43	46.04	2.49	1.41	46.79
Compost	low	R1	2.45	1.46	44.91	2.51	1.44	45.66	2.71	1.46	44.91
	high	R2	2.43	1.45	45.28	2.43	1.42	46.42	2.53	1.30	50.94
Mean			2.44	1.455	45.09	2.47	1.43	46.04	2.62	1.39	47.90
MNRM	low	R1	2.40	1.44	45.66	2.40	1.44	45.66	2.41	1.45	45.28
	high	R2	2.38	1.43	46.00	2.36	1.40	47.17	2.37	1.35	49.06
Mean			2.39	1.435	45.85	2.38	1.42	46.42	2.39	1.40	47.17

Their mixtures (1:1:1)	low	R1	2.38	1.41	46.79	2.48	1.42	46.42	2.45	1.36	48.68
	high	R2	2.32	1.34	49.43	2.34	1.31	50.57	2.39	1.32	50.19
Mean			2.35	1.375	48.11	2.41	1.365	48.50	2.42	1.34	49.43
Average	low	R1	2.41	1.430	46.04	2.47	1.437	45.77	2.53	1.423	46.30
	high	R2	2.38	1.423	45.42	2.44	1.385	47.74	2.43	1.343	49.32
Overall mean			2.395	1.426	46.33	2.455	1.411	46.75	2.48	1.383	47.81

Notes: 1-Each value is a mean of three replications and the obtained results were calculated on oven dry weight basis at 105 C° for 24 hours

2- **SHC** : Soil hydraulic conductivity ; **D_b** : Soil bulk density and **ρ_t** : Soil porosity in volume percent .

Table (9.2): Saturated hydraulic conductivity, bulk density and total porosity of studied soils after maize crop harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes.

Soil conditioner types and their mixtures	Conditioner application rates (w/w)	Soil moisture depletion levels from its available water capacity(AWSMD-levels)															
		Wet - treatment (30 % AWSMD) Light irrigation Short – intervals (every day)			Medium - treatment (50 % AWSMD) Moderate irrigation Median – intervals (3days)			Dry -treatment (70 % AWSMD) Heavy irrigation Long-intervals (8 days)									
		15.33 %SMC			13.5 % SMC			11.7 % SMC									
		Physical parameters															
SHC m/day			D _b Mg /m ³			ρ _t %			SHC m/day			D _b Mg /m ³			ρ _t %		
Control	Without additions	2.63	1.50	43.40	2.60	1.48	44.15	2.55	1.52	42.64							

Bentonite	low	R1	2.60	1.41	46.79	2.84	1.35	49.06	2.55	1.42	46.41
	high	R2	2.32	1.33	49.80	2.31	1.30	50.94	2.40	1.22	53.96
Mean			2.46	1.364	48.34	2.575	1.33	49.81	2.475	1.32	50.19
Compost	low	R1	2.80	1.38	47.92	3.00	1.35	49.57	3.95	1.35	49.06
	high	R2	2.42	1.35	49.06	2.32	1.30	50.94	3.17	1.28	51.70
Mean			2.61	1.365	48.50	2.66	1.325	50.00	3.56	1.315	50.38
MNRM	low	R1	2.64	1.40	47.17	2.50	1.28	51.70	2.86	1.38	47.92
	high	R2	2.20	1.34	49.43	2.48	1.40	47.17	3.10	1.30	50.94
Mean			2.42	1.37	48.30	2.49	1.34	49.44	2.98	1.34	49.43
Their mixtures (1:1:1)	low	R1	2.44	1.39	47.54	2.38	1.38	47.92	2.30	1.37	48.30
	high	R2	2.34	1.37	48.30	2.35	1.35	49.06	2.00	1.32	50.19
Mean			2.39	1.38	47.92	2.365	1.365	48.40	2.15	1.35	49.06
Average	low	R1	2.62	1.39	47.55	2.68	1.34	49.34	2.915	1.38	47.92
	high	R2	2.32	1.35	48.67	2.366	1.344	49.83	2.660	1.28	51.70
Overall mean			2.47	1.371	48.26	2.523	1.342	49.36	2.79	1.33	49.77

Notes:1-Each value is a mean of three replications and the obtained results were calculated on oven dry weight basis at 105 C° for 24 hours.

2- SHC : Saturated hydraulic conductivity ; D_b : Soil bulk density and ρ_t : Total porosity(in volume percent).

Table(10.1): Field capacity, permanent wilting point and available water capacity of studied soils after wheat crop harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes.

Soil conditioner types and their mixtures	Conditioner application rates (w/w)		Soil moisture depletion levels from its available water capacity(AWSMD-levels)								
			Wet- treatment (30 % AWSMD) Light irrigation Short – intervals (3 days)			Medium - treatment (50 % AWSMD) Moderate irrigation Median – intervals (6 days)			Dry - treatment (70 % AWSMD) Heavy irrigation Long- intervals (9 days)		
			14.45 % SMC			12.75 % SMC			11.08 % SMC		
			Soil moisture constants								
		SFC%	PWP %	AWC %	SFC%	PWP %	AWC%	SFC%	PWP %	AWC%	
Control	Without additions		17.0	8.5	8.5	16.6	8.2	8.4	16.4	8.2	8.2
Bentonite	Low	R1	17.7	8.7	9.0	17.4	8.6	8.8	16.8	8.4	8.4
	High	R2	18.4	9.0	9.4	17.8	8.8	9.0	17.3	8.5	8.8
Mean			18.05	8.85	9.2	17.6	8.7	8.9	17.05	8.45	8.6
Compost	Low	R1	17.7	8.8	8.9	17.3	8.5	8.8	17.2	8.5	8.7
	High	R2	18.1	8.8	9.3	18.5	8.9	9.6	17.5	8.8	8.7
Mean			17.9	8.8	9.1	17.9	8.7	9.2	17.35	8.65	8.7
MNRM	Low	R1	19.2	9.5	9.7	18.5	9.2	9.3	17.9	8.9	9.0
	High	R2	19.5	9.8	9.7	18.7	9.3	9.4	18.2	9.0	9.2

Mean			19.35	9.65	9.7	18.6	9.25	9.35	18.05	8.95	9.1
Their mixtures (1:1:1)	Low	R1	19.6	9.8	9.8	18.8	9.4	9.4	18.5	9.2	9.3
	High	R2	20.4	10.0	10.4	19.4	9.7	9.7	19.0	9.4	9.6
Mean			20.0	9.9	10.1	19.1	9.55	9.55	18.75	9.3	9.45
Average	Low	R1	18.55	9.2	9.35	18.0	8.93	9.075	17.6	8.75	8.85
	High	R2	19.10	9.4	9.70	18.6	9.18	9.425	18.0	8.93	9.07
Overall mean			18.83	9.3	9.53	18.03	9.05	9.25	17.8	8.84	8.96

Notes: 1-Each value is a mean of three replications and the obtained results were calculated on oven dry weight basis at 105 C° for 24 hours

2- SFC : soil field capacity ; PWP : permanent wilting point and AWC : soil available water capacity

Table(10.2): Field capacity, permanent wilting point and available water capacity of studied soils after maize crop harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes.

Soil conditioner types and their mixtures	Conditioner application rates (w/w)	Soil moisture depletion levels from its available water capacity(AWSMD-levels)		
		Wet- treatment (30 % AWSMD) Light irrigation Short – intervals (every day)	Medium - treatment (50 % AWSMD) Moderate irrigation Median – intervals (3 days)	Dry - treatment (70 % AWSMD) Heavy irrigation Long- intervals (8 days)
		15.33 %SMC	13.5 % SMC	11.7 % SMC

			Soil moisture constants								
			SFC %	PWP %	AWC %	SFC %	PWP %	AWC %	SFC %	PWP %	AWC %
Control	Without additions		18.0	9.0	9.0	17.8	9.2	8.6	17.9	8.8	9.1
Bentonite	Low	R1	18.5	9.3	9.2	18.3	9.5	8.8	19.2	9.0	10.2
	High	R2	18.9	9.5	9.4	18.8	9.6	9.2	19.8	9.2	10.6
Mean			18.7	9.4	9.3	18.55	9.55	9.0	19.5	9.1	10.4
Compost	Low	R1	19.5	9.8	9.7	19.6	10.0	9.6	20.3	10.1	10.2
	High	R2	19.7	10.0	9.7	20.3	10.4	9.9	20.5	10.3	10.2
Mean			19.6	9.9	9.7	19.95	10.2	9.75	20.4	10.2	10.2
MNRM	Low	R1	18.6	9.5	9.1	18.7	9.6	9.1	18.9	9.3	9.6
	High	R2	18.9	9.6	9.3	18.9	9.8	9.1	19.3	9.5	9.8
Mean			18.75	9.55	9.2	18.8	9.7	9.1	19.1	9.4	9.7
Their mixtures (1:1:1)	Low	R1	19.4	9.7	9.7	19.5	9.9	9.6	19.8	9.8	10.0
	High	R2	19.5	9.8	9.7	19.7	10.0	9.7	20.2	10.2	10.0
Mean			19.45	9.75	9.7	19.6	9.95	9.65	20.0	10.0	10.0
Average	Low	R1	19.0	9.58	9.43	19.02	9.75	9.275	19.55	9.55	10.0
	High	R2	19.25	9.73	9.53	19.43	9.95	9.476	19.95	9.8	10.15
Overall mean			19.12	9.65	9.47	19.23	9.85	9.375	19.75	9.67	10.07

Notes : 1-Each value is a mean of three replications and the obtained results are calculated on oven dry weight basis at 105 C° for 24 hours

2- SFC : soil field capacity ; PWP : permanent wilting point and AWC : soil available water capacity

Table(11.1): Phyto-availability (concentration) of soil macro-nutrients after wheat and maize crops harvesting as affected by natural soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes.

Soil conditioner types and their mixtures	Conditioner application rates (w/w)		After pot-wheat crop cultivation						After field-maize crop cultivation						
			Wet-treatment (30 % AWSMD) Light irrigation Short – intervals (3 days)		Medium -treatment (50 % AWSMD) Moderate irrigation Median – intervals (6 days)		Dry -treatment (70 % AWSMD) Heavy irrigation Long- intervals (9 days)		Wet-treatment (30 % AWSMD) Light irrigation Short – intervals (every day)		Medium -treatment (50 % AWSMD) Moderate irrigation Median – intervals (3 days)		Dry -treatment (70 % AWSMD) Heavy irrigation Long- intervals (8 days)		
			14.45 % SMC		12.75 % SMC		11.08 % SMC		15.33 % SMC		13.5 % SMC		11.7 % SMC		
			Soil nutritional status						Soil nutritional status						
Soil macro-nutrients phytoavailability ($mg\ kg^{-1}$ soil)															
				N	P	N	P	N	P	N	P	N	P	N	P
Control	Without additions		25.0	7.5	26.7	8.5	23.8	6.0	22.0	9.9	23.0	10.9	21.0	8.9	
Bentonite	low	R1	30.5	8.5	32.4	9.5	31.0	7.5	22.5	10.8	23.5	11.8	21.5	9.8	
	high	R2	35.6	9.5	35.5	10.5	31.3	8.4	25.5	11.5	26.5	12.5	24.5	10.5	
Mean			33.05	9.0	33.95	10.0	31.15	7.95	24.0	11.15	25.0	12.15	23.0	10.15	
Compost	low	R1	38.5	8.7	34.8	9.7	35.0	7.8	35.5	13.1	36.5	14.1	34.5	12.1	

	high	R2	39.9	9.5	39.8	10.9	36.5	8.5	40.8	13.5	41.8	14.5	39.8	12.5
Mean			39.2	9.1	37.3	10.3	35.75	8.15	38.15	13.3	39.15	14.3	37.15	12.3
MNRM	low	R1	37.8	10.2	46.5	11.2	37.2	9.2	30.2	12.5	31.2	11.5	29.2	11.5
	high	R2	40.5	10.5	47.3	11.5	40.0	9.9	35.3	12.8	36.3	13.8	34.3	11.8
Mean			39.15	10.35	46.9	11.35	38.6	9.55	32.75	12.65	33.75	12.65	31.75	11.65
Their mixtures (1:1:1)	low	R1	43.5	11.2	49.3	12.2	43.0	10.2	36.3	12.9	37.3	13.9	35.3	11.9
	high	R2	46.5	13.9	49.5	14.9	44.5	12.9	39.5	13.2	40.5	14.2	38.5	12.2
Mean			45.0	12.55	49.4	13.55	43.75	11.55	37.9	13.05	38.9	14.05	36.9	12.05
Average	low	R1	37.58	9.65	40.75	10.65	36.55	8.67	31.13	12.33	32.13	12.83	30.13	11.33
	high	R2	40.63	10.85	43.03	11.95	38.08	9.93	35.28	12.75	36.28	13.75	34.28	11.75
Overall mean			39.10	10.25	41.89	11.30	37.31	9.30	33.21	12.54	34.21	13.29	32.21	11.54

Notes : 1-Each value is a mean of 3 replications and all obtained values were calculated on oven dry weight basis at 105 C° for 24 hours

2-Each experimental plastic pot received (w/w) 120.6 kg-N fed⁻¹ (1.206 g- N pot⁻¹) as ammonium nitrate (33.5 % N); 13.54 kg-P fed⁻¹(135.4 mg-P pot⁻¹)as normal super phosphate (15.5 % P₂O₅) and 28.8 kg-K fed⁻¹(199.15 mg-K pot⁻¹) as potassium sulfate (48 % K₂O).

3- Each experimental plot area received 260 kg fed⁻¹ urea (46.5% N) equivalent 0.127 kg – N plot⁻¹ ; 200 kg fed⁻¹ ordinary super phosphate (15.5 % P₂O₅) equivalent (14.16 g-P plot⁻¹) and 50 kg fed⁻¹ potassium sulfate (48 % K₂O) equivalent 20.84 g-K plot⁻¹.

Table(12.1): Biomass grains and straw yields of wheat and maize crops after full maturity as affected by natural soil conditioner types, their mixtures and application rates under wet-treatment (light irrigation) .

Soil conditioner types and their mixtures	Conditioner application rates (w/w)		Biomass wheat crop yield						Biomass maize crop yield					
			Light irrigation (short – intervals 3 days)						Light irrigation (short – intervals every day)					
			Biomass grains yield		1000 grains weight (gm)	Harvest index (%)	Biomass straw (tepn)yield		Biomass grains yield		100 grains weight (gm)	Harvest index (%)	Biomass straw yield	
			g pot ⁻¹	(kg fed ⁻¹)			g pot ⁻¹	(kg fed ⁻¹)	(kg plot ⁻¹)	(kg fed ⁻¹)			(kg plot ⁻¹)	(kg fed ⁻¹)
Control	Without additions		13.77	1276.5	30.5	30.75	31.01	2875	1.950	1820	40.6	47	2.199	2053
Bentonite	low	R1	17.55	1627.5	40.5	34.14	29.84	2767	2.904	2710	42.5	49.18	2.999	2800
	high	R2	20.19	1872.0	40.8	40.35	33.87	3140	2.942	2746	42.8	47.72	3.219	3005
Mean			18.87	1749.8	40.65	37.25	31.86	2953	2.923	2728	42.65	48.45	3.110	2903
Compost	low	R1	15.81	1465.5	40.2	33.29	30.63	2840	3.129	2920	44.7	45.15	3.800	3547
	high	R2	18.72	1735.5	40.5	37.93	31.68	2937	3.185	2973	45.0	44.39	3.989	3724
Mean			17.26	1600.5	40.35	35.61	31.16	2888	3.157	2946	44.85	44.77	3.895	3635
MNRM	low	R1	19.66	1822.5	42.6	36.75	33.36	3093	3.054	2850	43.1	50.44	2.999	2800
	high	R2	23.93	2218.5	43.5	41.76	33.84	3137	3.107	2900	43.8	47.94	3.299	3080
Mean			21.79	2020.6	43.05	39.26	33.6	3115	3.080	2875	43.45	49.19	3.182	2970
Their mixtures (1:1:1)	low	R1	23.67	2194.5	46.2	38.64	36.35	3370	3.111	2903	43.7	47.05	3.500	3267
	high	R2	24.19	2242.6	44.5	40.30	37.07	3437	3.161	2950	44.5	44.89	3.879	3621
Mean			23.93	2218.6	45.35	39.47	36.71	3403	3.136	2926	44.10	45.94	3.690	3444
Average	low	R1	19.17	1777.0	42.37	35.89	32.54	3017	3.049	2846	43.5	47.95	3.325	3103

	high	R2	21.75	2017.0	42.32	40.17	34.11	3162	3.099	2892	44.0	46.23	3.697	3357
Overall mean			20.46	1897.3	42.35	38.05	33.33	3090	3.074	2869	43.75	47.09	3.461	3230

- Notes:** 1- Each figure is a mean of 3 variables and all obtained data were calculated on oven dry weight basis at 70 C° for 18 hours
 2- Ardeb of wheat grains = 150 kg and heml straw (tepn) = 250 kg, while, ardeb of maize grains = 140 kg
 3- Winter growing season elongated 135 days for wheat and growing summer season period elongated 93 days.
 4- Wet – treatment (light irrigation): 30 % AWSMD (short – intervals)(3 days) for wheat and every day for maize crop.

Table(12.2): Biomass grains and straw yields of wheat and maize crops after full maturity as affected by natural soil conditioner types, their mixtures and application rates under medium-treatment (moderate irrigation).

Soil conditioner types and their mixtures	Conditioner application rates (w/w)		Biomass wheat crop yield					Biomass maize crop yield						
			Moderate irrigation (median – intervals 6 days)					Moderate irrigation (median – intervals 3 days)						
			Biomass grains yield		1000 grains weight (gm)	Harvest Index (%)	Biomass straw (tepn)yield		Biomass grains yield		100 grains weight (gm)	Harvest Index (%)	Biomass straw yield	
			g pot ⁻¹	(kg fed ⁻¹)			g pot ⁻¹	(kg fed ⁻¹)	(kg plot ⁻¹)	(kg fed ⁻¹)			(kg plot ⁻¹)	(kg fed ⁻¹)
Control	Without additions		14.04	1302	30.6	32.69	28.91	2680	2.118	1977	41.2	42.46	2.870	2676
Bentonite	low	R 1	17.22	1596	40.3	36.25	30.28	2807	3.003	2803	42.6	49.21	3.099	2893
	high	R 2	20.16	1869	40.6	37.89	33.04	3063	3.096	2890	43.5	49.17	3.200	2987
Mean			18.69	1732.5	40.4	37.07	31.66	2935	3.050	2846	43.05	49.19	3.150	2940
Compost	low	R 1	15.97	1480.5	40.0	34.81	29.91	2773	3.204	2990	42.2	44.48	3.999	3733

	high	R 2	18.07	1675.5	40.2	36.84	30.98	2878	3.304	3084	45.2	44.03	4.199	3920
Mean			17.02	1578	40.1	35.82	30.45	2823	3.254	3037	43.7	44.26	4.100	3827
MNRM	low	R 1	19.25	1785	42.2	38.09	31.28	2900	3.107	2900	43.9	49.11	3.219	3005
	high	R 2	22.81	2115	42.9	41.05	32.76	3037	3.204	2990	44.5	45.16	3.894	3631
Mean			21.03	1950	42.5	39.57	32.02	2969	3.155	2945	44.2	47.14	3.565	3318
Their mixtures (1:1:1)	low	R 1	19.85	1840.5	43.6	38.56	31.63	2933	3.242	3026	44.5	44.83	3.989	3724
	high	R 2	23.67	2194.5	45.2	40.85	34.27	3177	3.296	3076	45.0	44.89	4.100	3827
Mean			21.76	2017.5	44.4	39.77	32.95	3055	3.269	3051	44.75	44.86	4.045	3776
Average	low	R 1	18.07	1675.5	41.53	36.99	30.78	2853	3.139	2930	43.3	46.91	3.579	3338
	high	R 2	21.178	1963.5	42.23	39.26	32.76	3038	3.225	3010	44.6	45.81	3.848	3591
Overall mean			19.63	1819.5	41.88	38.19	31.76	2946	3.182	2970	43.93	46.36	3.713	3465

Notes: 1- Each figure is a mean of 3 variables and all obtained data were calculated on oven dry weight basis at 70 C° for 18 hours

2- Medium – treatment (moderate irrigation): 50 % AWSMD (median – intervals)(6 days) for wheat and 3 days for maize crop.

3- Harvest index (%) = [Biomass grains yield(g pot⁻¹) /Biological crop yield (g pot⁻¹)] x 100 on oven dry weight basis at 70 C° for 18 hours.

Table(12.3): Biomass grains and straw yields of wheat and maize crops after full maturity as affected by natural soil conditioner types, their mixtures and application rates under dry-treatment (heavy irrigation).

Soil conditioner types and	Conditioner application	Biomass wheat crop yield	Biomass maize crop yield
		Heavy irrigation (long – intervals 9 days)	Heavy irrigation (long – intervals 8 days)

their mixtures	rates (w/w)		Biomass grains yield					Biomass straw (tepn) yield		Biomass grains yield		100 grains weight (gm)	Harvest index (%)	Biomass straw yield	
			g pot ⁻¹		(kg fed ⁻¹)	1000 grains weight (gm)	Harvest index (%)	g pot ⁻¹		(kg plot ⁻¹)				(kg plot ⁻¹)	(kg fed ⁻¹)
			(kg fed ⁻¹)					(kg fed ⁻¹)		(kg fed ⁻¹)					
Control	Without additions		10.79	1000	30.3	34.45	20.53	1903	2.036	1900	40.0	49.22	2.099	1960	
Bentonite	low	R 1	13.09	1213	39.5	36.22	23.05	2137	2.818	2630	41.0	49.45	2.88	2688	
	high	R 2	18.38	1704	40.1	38.87	28.91	2630	2.807	2620	42.1	48.35	2.99	2800	
Mean			15.74	1458	39.8	37.72	25.98	2408	2.812	2625	41.55	48.89	2.939	2744	
Compost	low	R 1	12.26	1137	38.9	33.24	24.62	2283	2.982	2783	41.2	46.43	3.44	3211	
	high	R 2	15.92	1476	40.1	37.41	26.64	2470	3.107	2900	42.3	44.4	3.89	3631	
Mean			14.09	1306	39.5	35.47	25.63	2376	3.044	2841	41.74	45.42	3.665	3421	
MNRM	low	R 1	15.06	1396	42.9	39.52	23.05	2137	2.989	2790	42.7	49.92	2.99	2800	
	high	R 2	24.79	2298	44.5	47.67	27.21	2523	3.001	2801	42.8	48.39	3.20	2987	
Mean			19.93	1847	43.7	44.23	25.13	2330	2.995	2795	42.73	49.16	3.099	2893	
Their mixtures (1:1:1)	low	R 1	16.31	1512	42.5	35.79	29.26	2713	3.054	2850	42.5	47.83	3.33	3108	
	high	R 2	20.31	1882	44.2	40.06	30.38	2817	3.096	2890	44.0	46.59	3.54	3313	
Mean			18.30	1697	43.4	38.02	29.82	2765	3.075	2870	43.25	47.21	3.44	3211	
Average	low	R 1	14.18	1314	40.95	36.2	24.99	2317	2.961	2763	41.9	48.41	3.16	2951	
	high	R 2	19.86	1840	42.51	41.23	28.29	2610	3.003	2803	42.7	46.93	3.41	3182	

Overall mean	17.02	1577	41.61	38.98	26.64	2463	2.982	2783	42.32	47.67	3.285	3067
--------------	-------	------	-------	-------	-------	------	-------	------	-------	-------	-------	------

Notes :

- 1- Each experimental plastic pot received (w/w) 360 kg fed⁻¹ NH₄NO₃ (33.5 % N) equivalent 1.206 g-N pot⁻¹; 200 kg fed⁻¹ ordinary superphosphate (15.5 % P₂O₅) equivalent (135.4 mg -P pot⁻¹) and 50 kg fed⁻¹ K₂SO₄ (48 % K₂O) equivalent 199.15 mg-K pot⁻¹).
- 2- Each experimental plot area received 260 kg fed⁻¹ urea (46.5 % N) equivalent 0.127 kg-N plot⁻¹; 200 kg fed⁻¹ ordinary superphosphate (15.5 % P₂O₅) equivalent (14.16 g -P plot⁻¹) and 50 kg fed⁻¹ K₂SO₄ (48 % K₂O) equivalent 20.84 g-K plot⁻¹).
- 3- Each figure is a mean of 3 variables and all obtained data were calculated on oven dry weight basis at 70 C° for 18 hours.
- 4- Dry – treatment (heavy irrigation): 70 % AWSMD (long – intervals)(9 days) for wheat and 8 days for maize crop.

Table(12.4): Yield components of wheat crop after full maturity as affected by soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes over the growing winter season period 2010/2011.

Soil conditioner types and their mixtures	Conditioner application rates (w/w)	Soil moisture depletion levels from its available water capacity (AWSMD-levels)													
		Wet - treatment (30 % AWSMD) Light irrigation Short- intervals (3 days)				Medium- treatment (50 % AWSMD) Moderate irrigation Median - intervals (6 days)				Dry -treatment (70 % AWSMD) Heavy irrigation Long-intervals (9 days)					
		14.45 % SMC				12.75 SMC				11.08 % SMC					
		Plant height (cm)	No of tillers /spike	No of spikelets/ spike	Spike length(cm)	Panicle mean weight ton fed ⁻¹	Plant height (cm)	No of tillers /spike	No of spikelets /spike	Spike length(cm)	Panicle mean weight ton fed ⁻¹	Plant height (cm)	No of tillers /spike	No of spikelets/ spike	Spike length(cm)

Control	Without additions		79.5	2.2	17	10.8	1.0	77.5	3.3	17.2	10.7	1.32	72.5	3.2	16.8	10.5	1.19
Bentonite	low	R1	85.6	3.6	17.5	11.5	1.76	82.5	3.4	17.2	11.3	1.75	80.5	3.1	16.9	11.0	1.55
	high	R2	88.5	3.8	17.9	11.8	2.06	84.5	3.6	17.5	11.5	2.04	81.5	3.3	17.5	11.3	1.83
Mean			87.05	3.7	17.7	11.65	1.91	83.5	3.5	17.35	11.4	1.895	81.0	3.2	17.2	11.15	1.69
Compost	low	R1	82.4	3.3	17.2	11.2	1.64	80.5	3.2	17.2	11.0	1.61	78.5	2.9	17.0	10.9	1.53
	high	R2	84.5	3.6	17.5	11.5	1.85	82.5	3.4	17.3	11.3	1.77	79.9	3.1	17.1	11.2	1.70
Mean			83.45	3.45	17.35	11.35	1.745	81.5	3.3	17.25	11.15	1.69	79.2	3.0	17.05	11.05	1.615
MNRM	low	R1	88.5	4.2	18.5	11.8	1.65	87.5	3.9	18.2	11.5	1.71	83.5	3.2	18.0	10.9	1.67
	high	R2	90.5	4.5	18.8	12.3	1.90	89.5	4.2	18.5	12.1	1.87	85.5	3.8	18.2	11.8	1.80
Mean			89.5	4.35	18.65	12.05	1.775	88.5	4.05	18.35	11.8	1.79	84.5	3.5	18.1	11.35	1.735
Their mixtures (1:1:1)	low	R1	91.5	4.6	19.2	12.0	2.03	90.5	4.3	18.8	11.9	2.00	85.5	4.0	18.4	11.5	1.88
	high	R2	92.5	4.7	19.8	12.8	2.37	91.5	4.6	19.5	12.5	2.37	87.4	4.3	19.2	12.2	2.12
Mean			92.0	4.65	19.5	12.4	2.20	91.0	4.45	19.15	12.2	2.185	86.45	4.15	18.8	11.85	2.00
Average	low	R1	91.3	4.5	19.2	12.3	2.09	90.3	4.35	18.95	12.1	2.08	85.96	3.98	18.6	11.7	1.93
	high	R2	91.8	4.6	19.44	12.3	2.17	90.84	4.42	19.10	12.1	2.16	86.32	4.10	18.7	11.81	1.98
Overall mean			91.6	4.6	19.36	12.34	2.10	90.6	4.38	19.02	12.13	2.12	86.14	4.04	18.69	11.77	1.95

Notes: Each figure is a mean of 3 replicates.

Table(5.5): Yield components of maize crop after full maturity as affected by soil conditioner types, their mixtures and application rates under different soil moisture depletion regimes over the summer growing season period 2011.

Soil conditioner types and

Soil moisture depletion levels from its available water capacity (AWSMD-levels)

their mixtures	Conditioners application rates (w/w)		Wet - treatment (30 % AWSMD) light irrigation Short- intervals (every day)	Medium- treatment (50 % AWSMD) Moderate irrigation Median - intervals (3 days)	Dry -treatment (70 % AWSMD) Heavy irrigation Long-intervals (8 days)						
			15.33 %SMC			13.5 % SMC			11.7 % SMC		
			Ear weight (g)	Plant height (cm)	Ear length (cm)	Ear weight (g)	Plant height (cm)	Ear length (cm)	Ear weight (g)	Plant height (cm)	Ear length (cm)
Control	Without additions		251.5	180.0	14.5	252.0	182.5	15.5	250.0	177.0	13.8
Bentonite	low	R1	288.4	190.0	15.5	290.0	195.3	16.0	287.5	188.0	15.0
	high	R2	290.4	195.0	16.0	292.5	196.0	16.7	290.0	194.0	14.5
Mean			289.4	192.5	15.75	291.25	195.65	16.35	288.75	191.0	14.75
Compost	low	R1	340.8	204.0	19.0	342.5	208.0	20.0	340.0	203.0	19.0
	high	R2	355.9	206.0	20.0	360.0	210.0	22.0	352.7	205.0	19.5
Mean			348.35	205.0	19.5	351.25	209.0	21.0	346.35	204.0	19.25
MNRM	low	R1	310.3	195.5	16.6	315.0	196.0	17.5	309.7	194.9	16.0
	high	R2	322.2	197.5	18.5	325.0	200.0	19.0	320.0	195.5	18.0
Mean			316.25	196.5	17.55	320.0	198.0	18.25	314.85	195.2	17.0
Their mixtures (1:1:1)	low	R1	333.6	200.0	19.0	335.0	205.0	20.0	332.0	199.0	18.7
	high	R2	343.7	203.0	19.5	340.0	205.5	22.0	340.0	204.0	19.0
Mean			338.65	201.5	19.25	337.5	205.25	21.0	336.0	201.5	18.85

Average	low	R1	318.27	197.37	17.52	320.6	201.07	18.37	317.3	196.22	17.05
	high	R2	328.05	200.37	18.50	329.4	202.87	19.92	325.7	199.63	17.38
Overall mean			323.16	198.87	18.01	324.98	201.97	19.14	321.48	197.92	17.21

Notes: Each figure is a mean of 3 replicates.

Soil Characters	Obtained values	
	Site (1)* Pot-experiment	Site(2)** Field-experiment
Chemical analysis		
Soil pH (1:2.5 soil-water suspension)	7.36	7.40
Electrical conductivity, EC dSm^{-1} (Soil past extract) at 25 C°	2.60	3.75
Saturation percentage(S.P)	38.0	40.0
Total soluble salts(T.S.S) $mg\ kg^{-1}$ soil	632(0.063%)	960(0.096%)
Calcium carbonate (CaCO ₃) %	0.40	0.60
Total soluble ions(1:5 Soil-water extractions)		
Soluble cations		
Ca ⁺² $meq\ L^{-1}$	1.10	1.00
Mg ⁺² $meq\ L^{-1}$	1.46	1.20
Na ⁺ $meq\ L^{-1}$	1.50	3.70
K ⁺ $meq\ L^{-1}$	0.14	0.10
Soluble anions		
CO ₃ ⁼ $meq\ L^{-1}$	0.00	0.00
HCO ₃ ⁻ $meq\ L^{-1}$	2.30	1.50
CL ⁻ $meq\ L^{-1}$	1.40	2.00
SO ₄ ⁻² $meq\ L^{-1}$	0.30	2.50
EC, dSm^{-1} (1:5 soil-water extraction)	0.416	0.602
Ionic strength (I.S) $mmoles\ L^{-1}$	4.50	4.45
Sodium adsorption ratio(SAR)	1.327	3.53
Soluble sodium percentage(SSP) %	35.71	61.7
Physical analysis		
Particle size distribution (g/100g soil)		
Coarse sand fraction %	65.0	50.0
Fine sand fraction %	10.0	5.50
Silt fraction %	15.0	31.0
Clay fraction %	10.0	13.5
Soil texture class	Loamy sand	Loamy sand
Soil bulk density(Db) $Mg\ m^{-3}$	1.57	1.55
Soil particle density (Dp) $Mg\ m^{-3}$	2.66	2.66
Total porosity(pt) on volume basis %	34.21	41.73
Soil saturated hydraulic conductivity (S.H.C) $m\ day^{-1}$	2.52	2.65

385

386

387

388

372